

Berezin quantization of noncommutative projective varieties

Andreas Andersson

Email: fornjotnr@hotmail.com

Max Planck Institute for Mathematics in the Sciences. Inselstrasse 22, D-04103 Leipzig, Germany
Wollongong University, School of Mathematics and Applied Statistics, 2522 Wollongong, Australia

Mathematics Subject Classification 2010 Primary: 58B32; Secondary: 58B34, 46L89

Keywords: Berezin quantization, Toeplitz algebras, Cuntz–Pimsner algebras, compact quantum groups, generalized inductive limits, noncommutative geometry, subproduct systems, noncommutative random walks, Arveson’s conjecture.

February 16, 2017

Abstract

We use operator algebras and operator theory to obtain new result concerning Berezin quantization of compact Kähler manifolds. Our main tool is the notion of subproduct systems of finite-dimensional Hilbert spaces, which enables all involved objects, such as the Toeplitz operators, to be very conveniently expressed in terms of shift operators compressed to a subspace of full Fock space. This subspace is not required to be contained in the symmetric Fock space, so from finite-dimensional matrix algebras we can construct noncommutative manifolds with extra structure generalizing that of a projective variety endowed with a positive Hermitian line bundle and a canonical Kähler metric in the class of the line bundle. Even in the commutative setting these constructions are very fruitful. Firstly, we show that the algebra of smooth functions on any smooth projective variety can be quantized in a strong sense of inductive limits, as was previously only accomplished for homogeneous manifolds. In this way the Kähler manifold is recovered exactly from quantization and not just approximately. Secondly, we obtain a strict quantization also for singular varieties. Thirdly, we show that the Arveson conjecture is true in full generality for shift operators compressed to the subspace of symmetric Fock space associated with any homogeneous ideal. For noncommutative examples we consider homogeneous spaces for compact matrix quantum groups which generalize q -deformed projective spaces, and we show that these can be obtained as the cores of Cuntz–Pimsner algebras constructed solely from the representation theory of the quantum group. We also discuss interesting connections with noncommutative random walks.

Contents

1	Introduction	3
2	Subproduct systems	5
2.1	Basic properties	5
2.2	Toeplitz algebras	6
2.2.1	The Toeplitz core	6
2.2.2	The right shifts	7
2.2.3	Normal ordering	7
2.3	Cuntz–Pimsner algebras	7

3	Review of quantization of projective varieties	8
3.1	Berezin quantization with prequantum condition	8
3.2	Projectively induced quantization	10
3.3	The circle bundle	12
3.4	Singular varieties	13
4	Inductive limits	13
4.1	Relaxed definition of inductive limits	14
4.2	Inductive limits from subproduct systems	16
4.3	Cuntz–Pimsner algebras from inductive limits	18
4.4	Formulas for covariant symbols	18
4.5	Changing the inner products	20
4.6	The isometries	20
5	Projective limits	23
5.1	Relaxing the notion of projective limit	23
5.2	Projective system for subproduct systems	23
5.3	The state on $\mathcal{O}_{\mathfrak{H}}$	24
5.3.1	$\mathfrak{J}_{l,m}$ as adjoint of $\iota_{m,l}$	24
5.3.2	The limit state	25
5.4	Contravariant symbols	26
5.5	The asymptotic multiplication	28
5.6	The adjoint of the total Toeplitz map	28
5.7	$\mathcal{O}_{\mathfrak{H}}^{(0)}$ as a projective limit	30
5.8	Strict quantization	30
5.9	$\mathcal{O}_{\mathfrak{H}}$ assembled from projective limits	32
5.10	Commutative case and the Arveson conjecture	32
6	Application to compact matrix quantum groups	34
6.1	Compact matrix quantum groups	34
6.1.1	Representations and actions	34
6.1.2	Universal quantum groups	36
6.1.3	The dual discrete quantum group	36
6.2	First-row and first-column algebras	38
6.3	Subproduct systems of \mathbb{G} -representations	39
6.4	Berezin quantization of $C(\mathbb{G}/\mathbb{K})$	40
6.4.1	Covariant symbols as first-row matrix coefficients	40
6.4.2	Intertwining the actions	40
6.4.3	$C(\mathbb{G}/\mathbb{K})$ as an inductive limit	41
6.4.4	$C(\mathbb{S}_{\mathbb{G}})$ as an inductive limit	42
6.5	Comparison with Poisson and Martin boundaries	43
6.5.1	Poisson integral versus total Toeplitz map	43
6.5.2	Markov operator	43
6.5.3	Martin boundaries	44
7	Concluding remarks	45
8	References	46

1 Introduction

With motivations from physics, Berezin introduced a way of approximating certain compact Kähler manifolds M by finite-dimensional matrix algebras $\mathcal{B}(\mathfrak{H}_m)$ parameterized by $m \in \mathbb{N}_0$ [Bere1], [Bere2], [CGR1], [Sch1], [Lan2]. When $M = G/K$ is a homogeneous space of some Lie group G , each Hilbert space \mathfrak{H}_m carries an irreducible representation of G , and it is obtained by the Borel–Weil construction: \mathfrak{H}_m is the space of holomorphic sections of a suitable line bundle $L^{\otimes m}$ over M .

In order to apply Berezin quantization to quantum physics, the parameters (time or temperature or energy etc.) should be chosen such that the limit $m \rightarrow \infty$ simplifies the description of the system at hand. This is a powerful method; for instance it gives the Hartree–Fock approximation as a special case [Raj1]. The limit behavior is captured by a classical (compact Kähler) manifold M .

Nowadays it is however becoming more and more important to have a versatile theory of open quantum systems. Recently we observed how to obtain a simplifying infinite- m limit in the standard framework of quantum channels as driving the evolution of open quantum systems [An4, An5]. It so happens though, that the infinite- m system is (in general) not given by a classical manifold, but by a *noncommutative* manifold, i.e. there is a noncommutative algebra $C^\infty(\mathbb{M})$ which is supposed to encode the properties of the dynamics and which is a surprisingly good analogue of the commutative algebra $C^\infty(M)$ of smooth functions on a compact manifold M . Here we use the symbol \mathbb{M} for a nonexistent object defined by the algebra $C^\infty(\mathbb{M})$, while M denotes an honest manifold.

The “noncommutative Kähler manifolds” \mathbb{M} appearing in this way generalize only a special kind of compact Kähler manifolds, namely (complex) projective manifolds $M \subset \mathbb{CP}^{n-1}$ (also singular varieties can occur though). The C^* -algebras $C(\mathbb{M})$ defining these noncommutative manifolds will be constructed in such a way that they possess a lot of extra structure generalizing that complex-analytic structure, a positive line bundle and a Kähler metric.

In the classical setting of Berezin quantization of smooth projective varieties, our main new result is that we can obtain $C^\infty(M)$ operator-algebraically as an inductive limit. In this way the algebraic structure of $C^\infty(M)$ is not merely approximated by finite-dimensional matrix algebras via Toeplitz operators, but also via covariant symbols and we can take an actual limit to recover $C^\infty(M)$ completely, together with a canonical Kähler metric on M obtained as an inductive limit of tracial states. These constructions work only when the “Berezin transforms” converge to the identity map on $C^\infty(M)$, and our result relies on an idea of how this can be accomplished without assuming that the projective manifold M is “balanced”. We refer to §3.2 for a detailed discussion of the commutative setting.

Smooth projective varieties include in particular all coadjoint orbits G/K equipped with the Kirillov Kähler form and, going noncommutative, we observe that every compact matrix quantum group \mathbb{G} defines such a “manifold” \mathbb{G}/\mathbb{K} . A “noncommutative projective variety” will be given by a sequence $\mathfrak{H}_\bullet = (\mathfrak{H}_m)_{m \in \mathbb{N}_0}$ of finite-dimensional Hilbert spaces such that $\mathfrak{H}_{m+l} \subseteq \mathfrak{H}_m \otimes \mathfrak{H}_l$ for all $m, l \in \mathbb{N}_0$. Such a sequence has been referred to as a “subproduct system” [ShSo1] and it generalizes the structure needed to perform ordinary Berezin quantization. Our main aim is then to show that the C^* -algebra $C(\mathbb{G}/\mathbb{K})$ can be recovered from a suitably chosen \mathfrak{H}_\bullet via a noncommutative version of strict quantization.

In fact the definition of \mathbb{G}/\mathbb{K} is very simple. Let \mathbb{G} be a compact matrix quantum group with defining unitary representation $u \in M_n(\mathbb{C}) \otimes C(\mathbb{G})$. We let $z_j := u_{1,j}$ for $j = 1, \dots, n$ denote the elements of the first row of u . Then $C(\mathbb{G}/\mathbb{K})$ is defined as the C^* -algebra generated by elements of the form $z_{j_1} \cdots z_{j_m} z_{k_m}^* \cdots z_{k_1}^*$ for all multi-indices (j_1, \dots, j_m) and (k_1, \dots, k_m) of equal length. We also denote by $C(\mathbb{S}_{\mathbb{G}})$ the C^* -algebra generated by z_1, \dots, z_n (and refer to it as the “first-row algebra”).

We can summarize one of the main results of this paper in the following way. Suppose that the compact matrix quantum group \mathbb{G} is such that every element of $C(\mathbb{G}/\mathbb{K})$ can be “normally ordered”, in the sense that all z_j^* ’s are to the right of the z_k ’s (this is a crucial assumption). Assume also that the Haar state is faithful on $C(\mathbb{G}/\mathbb{K})$.

Theorem 1.1. *For each $m \in \mathbb{N}_0$, let \mathfrak{H}_m be the Hilbert space spanned by products $z_{j_1} \cdots z_{j_m}$ with inner product coming from the Haar measure on \mathbb{G} (the sequence of \mathfrak{H}_m ’s is the “ \mathbb{G} -subproduct*

system" \mathfrak{H}_\bullet). Then there are explicit injections ("Berezin covariant symbol maps")

$$\varsigma_{\mathbb{G}}^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow C(\mathbb{G}/\mathbb{K})$$

and explicit surjections ("Toeplitz quantization maps")

$$\zeta_{\mathbb{G}}^{(m)} : C(\mathbb{G}/\mathbb{K}) \rightarrow \mathcal{B}(\mathfrak{H}_m)$$

such that $\varsigma_{\mathbb{G}}^{(m)} \circ \zeta_{\mathbb{G}}^{(m)}$ converges point-norm to the identity on $C(\mathbb{G}/\mathbb{K})$ as $m \rightarrow \infty$. This effects an isomorphism

$$C(\mathbb{S}_{\mathbb{G}}) \cong \mathcal{O}_{\mathfrak{H}}$$

of the first-row algebra $C(\mathbb{S}_{\mathbb{G}})$ with the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ of the subproduct system \mathfrak{H}_\bullet , and this isomorphism is equivariant for the natural ergodic actions of \mathbb{G} and its discrete dual group $\hat{\mathbb{G}}$.

Denoting by $\mathcal{O}_{\mathfrak{H}}^{(0)}$ the $U(1)$ -invariant part of $\mathcal{O}_{\mathfrak{H}}$ under the gauge action, we also have

$$C(\mathbb{G}/\mathbb{K}) \cong \mathcal{O}_{\mathfrak{H}}^{(0)}, \quad (1.1)$$

and it is the C^* -algebra $\mathcal{O}_{\mathfrak{H}}^{(0)}$ which will occupy most of the paper. We will first realize $\mathcal{O}_{\mathfrak{H}}^{(0)}$ as a "generalized inductive limit" in the sense of [BK1], [Haw1], and then as a generalized projective limit in the spirit of [Haw1]. Then we do the same thing for $C(\mathbb{G}/\mathbb{K})$ to obtain the desired isomorphism (1.1).

A Berezin quantization for compact quantum groups with tracial Haar state was first considered in [Sain].

The idea of looking at the \mathbb{G} -subproduct system was partially motivated by Woronowicz' reconstruction of a compact matrix quantum group \mathbb{G} from its irreducible representations [Wor3]. Since \mathfrak{H}_\bullet only contains a subset of all irreducible representations (in general), we recover not $C(\mathbb{G})$ but $C(\mathbb{G}/\mathbb{K})$.

For a classical manifold M , with quantization defined by a line bundle L over M , elements of $\mathcal{O}_{\mathfrak{H}}^{(k)}$ are continuous sections of the line bundle $L^{\otimes k}$. The subspace $\mathfrak{H}_m \subset \mathcal{O}_{\mathfrak{H}}^{(m)}$ consists of the holomorphic sections of the line bundle. For $M = G/K$ a homogeneous space, the \mathfrak{H}_m 's are irreducible representations of G . In general, the subspaces $\mathcal{O}_{\mathfrak{H}}^{(k)}$ for $k \in \mathbb{Z} \setminus \{0\}$ are Hilbert modules over $\mathcal{O}_{\mathfrak{H}}^{(0)}$, and for a quantum homogeneous space $\mathbb{M} = \mathbb{G}/\mathbb{K}$ each \mathfrak{H}_m is an irreducible representation of \mathbb{G} . In this sense, $\mathcal{O}_{\mathfrak{H}}$ is a kind of "Borel–Weil algebra" for \mathbb{G} (cf. [Seg1, Thm. 14.1]).

We shall also compare our results with "noncommutative random walks" on duals of compact quantum groups [Iz1], [INT1], [Iz4]. The Toeplitz core $\mathcal{T}_{\mathfrak{H}}^{(0)}$ plays the role of Martin compactification of a walk restricted to the "dual" of \mathbb{G}/\mathbb{K} while $\mathcal{O}_{\mathfrak{H}}^{(0)}$ is the boundary of the walk. Recalling that the simplest Cuntz–Pimsner algebras $C(\mathbb{S}^{2n-1})$ of functions on spheres behave like boundaries of the corresponding Toeplitz algebras, these observations are not too surprising.

The inductive and projective limits mentioned above encode the data of a "strict quantization", as needed to generalize the classical setting, but they are so much more convenient than just knowing that there is a strict quantization. The notion of subproduct systems and the associated operator algebras of shift operators provide a machinery for explicit calculations that has previously been available mainly in the case of \mathbb{CP}^{n-1} , where Berezin quantization and fuzzy geometry has been successfully described in terms of creation and annihilation operators (the unnormalized shift operators) (see e.g. [BDLMC]). Finally, the identification of these shift-operator algebras with inductive limits allows us to solve the ten years-old Arveson conjecture.

Acknowledgment. The author thanks Adam Rennie for great comments, support and inspiration. Many thanks also to Orr Shalit and Guy Salomon for finding an important error in an earlier version of this paper. We thank Orr Shalit also for other remarks of great value.

2 Subproduct systems

2.1 Basic properties

In this paper we write $\mathbb{N}_0 := \mathbb{N} \cup \{0\} = \{0, 1, 2, \dots\}$.

Definition 2.1. [[ShSo1, Def. 6.2]] A **subproduct system** is a sequence $\mathfrak{H}_\bullet = (\mathfrak{H}_m)_{m \in \mathbb{N}_0}$ of finite-dimensional Hilbert space \mathfrak{H}_m such that $\mathfrak{H}_0 = \mathbb{C}$ and

$$\mathfrak{H}_{m+l} \subseteq \mathfrak{H}_m \otimes \mathfrak{H}_l, \quad \forall m, l \in \mathbb{N}_0. \quad (2.1)$$

We shall always denote \mathfrak{H}_1 by \mathfrak{H} and, throughout this paper, $n \in \mathbb{N}$ will always be the dimension of \mathfrak{H} ,

$$\mathfrak{H} \cong \mathbb{C}^n.$$

Example 2.2. Given a finite-dimensional Hilbert space \mathfrak{H} , we set $\mathfrak{H}_m := \mathfrak{H}^{\otimes m}$ for each m and refer to it as the “product system” associated with \mathfrak{H} . Another example of a subproduct system is obtained by taking $\mathfrak{H}_m := \mathfrak{H}^{\vee m}$ to be the m th symmetric power of \mathfrak{H} ; this is the “symmetric subproduct system”.

Definition 2.3. A subproduct system is **commutative** if $\mathfrak{H}_m \subseteq \mathfrak{H}^{\vee m}$ for all $m \in \mathbb{N}_0$.

Lemma 2.4. [[ShSo1, Lemma 6.1]] Let \mathfrak{H}_\bullet be a subproduct system. Then the projections $p_m : \mathfrak{H}^{\otimes m} \rightarrow \mathfrak{H}_m$ satisfy

$$p_l(p_m \otimes p_{l-m})p_l = p_l = p_l(p_{l-m} \otimes p_m)p_l \quad (2.2)$$

whenever $m \leq l$.

Proof. Replacing l by $l - m$ for $m \leq l$, the condition (2.1) reads

$$\mathfrak{H}_l \subseteq \mathfrak{H}_m \otimes \mathfrak{H}_{l-m}, \quad \forall m \leq l \in \mathbb{N}_0. \quad (2.3)$$

Writing (2.3) in terms of projections,

$$p_l \leq p_m \otimes p_{l-m},$$

the result is clear. \square

Definition 2.5. The **Fock space** associated with a subproduct system \mathfrak{H}_\bullet is the Hilbert space

$$\mathfrak{H}_{\mathbb{N}} := \bigoplus_{m \in \mathbb{N}_0} \mathfrak{H}_m.$$

We regard $\mathfrak{H}_{\mathbb{N}}$ as a subspace of full Fock space $\mathfrak{H}^{\otimes \mathbb{N}} := \bigoplus_{m \in \mathbb{N}_0} \mathfrak{H}^{\otimes m}$ and denote by $p_{\mathbb{N}} = \sum_{m \in \mathbb{N}_0} p_m$ the projection from $\mathfrak{H}^{\otimes \mathbb{N}}$ onto $\mathfrak{H}_{\mathbb{N}}$. Thus $p_{\mathbb{N}}$ is the identity in $\mathcal{B}(\mathfrak{H}_{\mathbb{N}})$, just as p_m is the identity in $\mathcal{B}(\mathfrak{H}_m)$.

Example 2.6. If $\mathfrak{H}_\bullet = \mathfrak{H}^{\otimes \bullet}$ is the product system over a fixed Hilbert space \mathfrak{H} then $\mathfrak{H}_{\mathbb{N}}$ is the full (or “Boltzmannian”) Fock space $\mathfrak{H}^{\otimes \mathbb{N}}$ over \mathfrak{H} .

Example 2.7. If $\mathfrak{H}_\bullet = \mathfrak{H}^{\vee \bullet}$ is the full commutative subproduct system then $\mathfrak{H}_{\mathbb{N}}$ is the symmetric (or “Bosonic”) Fock space $\mathfrak{H}^{\vee \mathbb{N}} := \bigoplus_{m \in \mathbb{N}_0} \mathfrak{H}^{\vee m}$ over \mathfrak{H} .

Notation 2.8. We denote by $\mathbb{C}\langle \mathbf{z} \rangle = \mathbb{C}\langle z_1, \dots, z_n \rangle$ the algebra of polynomials in n freely commuting variables. We denote by $\mathbb{C}[\mathbf{z}] = \mathbb{C}[z_1, \dots, z_n]$ the algebra of polynomials in n commuting variables.

For any polynomial $f(z_1, \dots, z_n) = \sum_{j_1, \dots, j_n} f_{j_1, \dots, j_n} z_1^{j_1} \cdots z_n^{j_n}$ in $\mathbb{C}\langle z_1, \dots, z_n \rangle$, evaluation on the basis e_1, \dots, e_n for $\mathfrak{H} = \mathfrak{H}_1$ defines an element in Fock space,

$$f(e_1, \dots, e_n) := \sum_{j_1, \dots, j_n} f_{j_1, \dots, j_n} e_1^{j_1} \otimes \cdots \otimes e_n^{j_n} \in \mathfrak{H}_{\mathbb{N}},$$

and f is homogeneous iff

$$f(e_1, \dots, e_n) \in \mathfrak{H}^{\otimes m}, \quad \text{for some } m \in \mathbb{N}_0.$$

Lemma 2.9. *Every subproduct system $\mathfrak{H}_\bullet = (\mathfrak{H}_m)_{m \in \mathbb{N}_0}$ defines a homogeneous ideal in $\mathbb{C}\langle \mathbf{z} \rangle$, where $n = \dim(\mathfrak{H})$. Conversely, every homogeneous ideal in $\mathbb{C}\langle \mathbf{z} \rangle$ corresponds to a subproduct system [ShSo1, Prop. 7.2]. For commutative subproduct systems the same is true with $\mathbb{C}[\mathbf{z}]$ instead [DRS1, §2.3].*

Sketch of proof. Given \mathfrak{H}_\bullet , define a homogeneous ideal \mathcal{I} in $\mathbb{C}\langle \mathbf{z} \rangle$ by

$$\mathcal{I} := \{f \in \mathbb{C}\langle \mathbf{z} \rangle \mid f(e_1, \dots, e_n) \in \mathfrak{H}^{\otimes m} \ominus \mathfrak{H}_m \text{ for some } m \in \mathbb{N}_0\}.$$

Conversely, given a homogeneous ideal \mathcal{I} , we associate the Hilbert spaces

$$\mathfrak{H}_m := \mathfrak{H}^{\otimes m} \ominus \{f(e_1, \dots, e_n) \mid f \in \mathcal{I}^{(m)}\},$$

where $\mathcal{I}^{(m)}$ is the degree- m component of \mathcal{I} .

2.2 Toeplitz algebras

Having fixed a subproduct system \mathfrak{H}_\bullet , we shall always denote by S_1, \dots, S_n the operators on Fock space $\mathfrak{H}_\mathbb{N}$ defined by

$$S_k \phi := p_{m+1}(e_k \otimes \phi), \quad \forall \phi \in \mathfrak{H}_m$$

for all $m \in \mathbb{N}_0$, where $p_{m+1} : \mathfrak{H}^{\otimes(m+1)} \rightarrow \mathfrak{H}_{m+1}$ is the orthogonal projection. They are the compressions to $\mathfrak{H}_\mathbb{N}$ of the left shifts $\psi \rightarrow e_k \otimes \psi$ on full Fock space $\mathfrak{H}^{\otimes \mathbb{N}}$. For more about compressed n -tuples of shift operators, see [Pop1], [Pop2], [ShSo1], [DRS1], [DRS2].

Definition 2.10. The **Toeplitz algebra** of a subproduct system \mathfrak{H}_\bullet is the unital C^* -algebra $\mathcal{T}_{\mathfrak{H}}$ of operators on $\mathfrak{H}_\mathbb{N}$ generated by the shifts S_1, \dots, S_n .

The adjoint S_k^* of S_k preserves the subspace $\mathfrak{H}_\mathbb{N}$ so S_k^* is just the restriction to $\mathfrak{H}_\mathbb{N}$ of the backward shift on $\mathfrak{H}^{\otimes \mathbb{N}}$.

Definition 2.11. The **vacuum state** on the Toeplitz algebra $\mathcal{T}_{\mathfrak{H}}$ is the restriction $\hat{\varepsilon} : \mathcal{T}_{\mathfrak{H}} \rightarrow \mathbb{C}$ of the vector state on $\mathcal{B}(\mathfrak{H}_\mathbb{N})$ defined by the unit vector $\Omega \in \mathfrak{H}_0 = \mathbb{C}$. That is,

$$\hat{\varepsilon}(X) := \langle \Omega | X \Omega \rangle, \quad \forall X \in \mathcal{B}(\mathfrak{H}_\mathbb{N}).$$

If p_0 denotes the unit in $\mathcal{B}(\mathfrak{H}_0)$ then

$$X p_0 = \hat{\varepsilon}(X) p_0 = p_0 X, \quad \forall X \in \mathcal{B}(\mathfrak{H}_\mathbb{N}). \quad (2.4)$$

Notation 2.12. Let \mathbb{F}_n^+ be the free unital semigroup generated by n elements $1, \dots, n$ (the empty word \emptyset is the identity in \mathbb{F}_n^+). We write a word $\mathbf{k} \in \mathbb{F}_n^+$ as $\mathbf{k} = k_1 \cdots k_m$ and refer to $|\mathbf{k}| := m$ as the **length** of \mathbf{k} . For the shifts S_1, \dots, S_n and the basis vectors e_1, \dots, e_n we then write

$$S_{\mathbf{k}} := S_{k_1} \cdots S_{k_m}, \quad S_{\mathbf{k}}^* := (S_{\mathbf{k}})^* = S_{k_m}^* \cdots S_{k_1}^*,$$

$$e_{\mathbf{k}} := e_{k_1} \otimes \cdots \otimes e_{k_m},$$

and similarly for other n -tuples of elements defined below. Finally, $\mathbf{j}\mathbf{k} := j_1 \cdots j_l k_1 \cdots k_m$ for $\mathbf{j}, \mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{j}| = l$ and $|\mathbf{k}| = m$.

2.2.1 The Toeplitz core

Let $N = \bigoplus_m m p_m$ be the **number operator** on $\mathfrak{H}_\mathbb{N}$. It generates a unitary group on $\mathfrak{H}_\mathbb{N}$ which implements an action γ_\bullet of the circle group $U(1)$ on $\mathcal{T}_{\mathfrak{H}}$,

$$\gamma_t(S_k) := e^{it} S_k, \quad \forall e^{it} \in U(1), k \in \{1, \dots, n\}, \quad (2.5)$$

referred to as the **gauge action** on $\mathcal{T}_{\mathfrak{H}}$.

The gauge action (2.5) splits $\mathcal{T}_{\mathfrak{H}}$ into the C^* -direct sum of the subspaces

$$\mathcal{T}_{\mathfrak{H}}^{(k)} := \{T \in \mathcal{T}_{\mathfrak{H}} \mid \gamma_t(T) = e^{ikt} T \text{ for all } e^{it} \in U(1)\}, \quad k \in \mathbb{Z}.$$

The fixed-point subalgebra $\mathcal{T}_{\mathfrak{H}}^{(0)}$ (the **Toeplitz core**) will be of great importance to us. It is generated by polynomials in the shifts S_j and S_k^* which are “homogeneous of degree zero” in the sense that each term contains equally many forward shifts S_j as backward shifts S_k^* .

2.2.2 The right shifts

In addition to the “sinister” shift S_k by the basis vector $e_k \in \mathfrak{H}$, we shall need the “rectus” shift

$$R_k \psi := p_{m+1}(\psi \otimes e_k), \quad \forall \psi \in \mathfrak{H}_m, m \in \mathbb{N}_0 \quad (2.6)$$

acting on the same Fock space $\mathfrak{H}_{\mathbb{N}}$. Note that R_k commutes with each S_j , and that R_k^* commutes with each S_j^* , but $[R_j, S_k^*] \neq 0$.

The following formulas will be used extensively.

Lemma 2.13. *For all $m, l \in \mathbb{N}$ with $m \leq l$ we have*

$$\sum_{|\mathbf{r}|=m} S_{\mathbf{r}} S_{\mathbf{r}}^*|_{\mathfrak{H}_l} = p_l = \sum_{|\mathbf{r}|=m} R_{\mathbf{r}} R_{\mathbf{r}}^*|_{\mathfrak{H}_l}.$$

In particular,

$$\sum_{|\mathbf{r}|=m} S_{\mathbf{r}} S_{\mathbf{r}}^* = p_{\mathbb{N}} - |\Omega\rangle\langle\Omega|. \quad (2.7)$$

Proof. We have $S_{\mathbf{r}} S_{\mathbf{s}}^*|_{\mathfrak{H}_l} = p_l(|e_{\mathbf{r}}\rangle\langle e_{\mathbf{s}}| \otimes p_{l-m})p_l = p_l(|e_{\mathbf{r}}\rangle\langle e_{\mathbf{s}}| \otimes p_{l-m})p_l$, so

$$\begin{aligned} \sum_{|\mathbf{r}|=m} S_{\mathbf{r}} S_{\mathbf{r}}^*|_{\mathfrak{H}_l} &= p_l \left(\sum_{|\mathbf{r}|=m} |e_{\mathbf{r}}\rangle\langle e_{\mathbf{r}}| \otimes p_{l-m} \right) p_l \\ &= p_l(p_m \otimes p_{l-m})p_l = p_l, \end{aligned}$$

and similarly for the right shifts. \square

From (2.7) we see that the vacuum projection $p_0 = |\Omega\rangle\langle\Omega|$ belongs to the Toeplitz algebra. Considering multiplying p_0 from both sides with different shift operators, it is a simple matter to deduce the following.

Corollary 2.14. *The Toeplitz algebra $\mathcal{T}_{\mathfrak{H}}$ contains the C^* -algebra \mathcal{K} of all compact operators on Fock space $\mathfrak{H}_{\mathbb{N}}$ as a norm-closed two-sided ideal.*

2.2.3 Normal ordering

In the present paper we will obtain some basic but results about the Toeplitz core, which we collect here for convenience.

Lemma 2.15 (Normal ordering). *Let \mathfrak{H}_{\bullet} be a subproduct system. Let $\mathcal{A}_{\mathfrak{H}}$ denote the norm-closed (non-*) algebra generated by the shifts S_1, \dots, S_n and the identity in $\mathcal{B}(\mathfrak{H}_{\mathbb{N}})$. Then*

- (i) $\overline{\text{span}}(\mathcal{A}_{\mathfrak{H}} \mathcal{A}_{\mathfrak{H}}^*)$ is an algebra,
- (ii) $\mathcal{T}_{\mathfrak{H}} = \overline{\text{span}}(\mathcal{A}_{\mathfrak{H}} \mathcal{A}_{\mathfrak{H}}^*)$, and
- (iii) $\mathcal{T}_{\mathfrak{H}} = \overline{\text{span}}(\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}} \cup \mathcal{K})$.

The proof of (i) and (ii) will be given in Lemma 4.10 and (iii) will be obtained in §5.7. For the moment we just note that the C^* -algebra \mathcal{K} of compact operator is contained in $\overline{\text{span}}(\mathcal{A}_{\mathfrak{H}} \mathcal{A}_{\mathfrak{H}}^*)$.

2.3 Cuntz–Pimsner algebras

Definition 2.16 ([Vis2, Cor. 3.2]). The **Cuntz–Pimsner algebra** of a subproduct system \mathfrak{H}_{\bullet} is the quotient of the Toeplitz algebra $\mathcal{T}_{\mathfrak{H}}$ by the ideal \mathcal{K} of all compact operators on $\mathfrak{H}_{\mathbb{N}}$,

$$\mathcal{O}_{\mathfrak{H}} := \mathcal{T}_{\mathfrak{H}} / \mathcal{K}.$$

We denote by Z_1, \dots, Z_n the generators of $\mathcal{O}_{\mathfrak{H}}$, i.e. the images of the shifts S_1, \dots, S_n in the quotient. They satisfy the **sphere relation**

$$\sum_{k=1}^n Z_k Z_k^* = \mathbf{1},$$

which suggests viewing $\mathcal{O}_{\mathfrak{H}}$ as the “boundary” of $\mathcal{T}_{\mathfrak{H}}$; in the latter holds $\sum_{k=1}^n S_k S_k^* \leq \mathbf{1}$, as we saw in (2.7).

The formula (2.5), but with Z_k replacing S_k , defines the gauge action on $\mathcal{O}_{\mathfrak{H}}$, which gives a splitting

$$\mathcal{O}_{\mathfrak{H}} = \overline{\bigoplus_{k \in \mathbb{Z}} \mathcal{O}_{\mathfrak{H}}^{(k)}}^{\|\cdot\|}$$

of $\mathcal{O}_{\mathfrak{H}}$ into spectral subspaces for this $U(1)$ -action.

Remark 2.17 (Known examples). The most straightforward example of a subproduct Cuntz–Pimsner algebra is the Cuntz algebra \mathcal{O}_n , obtained from $\mathfrak{H}_{\bullet} = \mathfrak{H}^{\otimes \bullet}$. As a commutative example, $\mathcal{O}_{\mathfrak{H}}$ for the symmetric subproduct system $\mathfrak{H}^{\vee \bullet}$ was shown in [Ar6] to be isomorphic to the C^* -algebra $C(\mathbb{S}^{2n-1})$ of continuous functions on the unit sphere $\mathbb{S}^{2n-1} \subset \mathbb{C}^n$. Cuntz–Pimsner algebras coming from monomial ideals were described in [KaSh1].

It was conjectured in [Ar8] that $\mathcal{O}_{\mathfrak{H}}$ is commutative for any commutative subproduct system.

For any subproduct system \mathfrak{H}_{\bullet} , the spectral subspaces $\mathcal{O}_{\mathfrak{H}}^{(k)}$ for the gauge action on $\mathcal{O}_{\mathfrak{H}}$ are Hilbert C^* -bimodules over the fixed-point subalgebra $\mathcal{O}_{\mathfrak{H}}^{(0)}$, with left and right inner products

$$\langle \xi | \eta \rangle_{\text{right}} := \xi^* \eta, \quad \langle \xi | \eta \rangle_{\text{left}} := \xi \eta^* \quad (2.8)$$

for $\xi, \eta \in \mathcal{O}_{\mathfrak{H}}^{(k)}$.

3 Review of quantization of projective varieties

Let us formulate Berezin quantization of complex submanifolds of projective n -space \mathbb{CP}^{n-1} in terms of subproduct systems, just to make it clear how the results of the subsequent sections relate to the classical ones.

3.1 Berezin quantization with prequantum condition

Recall that Chow’s theorem says that a submanifold of projective space $\mathbb{P}[\mathbb{C}^n] = \mathbb{CP}^{n-1}$ is a nonsingular (i.e. smooth) projective variety, i.e. the zero-set of some finitely generated homogeneous ideal in $\mathbb{C}[z_1, \dots, z_n]$. These manifolds can be characterized without even referring to \mathbb{CP}^{n-1} (see Lemma 3.3 below), but for this we need to recall some complex geometry. A **Kähler manifold** is a pair (M, ω) consisting of a complex manifold M and a closed nondegenerate 2-form ω on M which equals the imaginary part of a Hermitian metric on M .

Remark 3.1 (Poisson bracket). The Kähler form ω is in particular a symplectic (i.e. closed and nondegenerate) form, making M a symplectic manifold. The nondegeneracy of ω allows us to use the inverse ω^{-1} to define a Poisson bracket on $C^\infty(M)$ by

$$\{f, g\} := \omega^{j,k} \frac{\partial f}{\partial x_j} \frac{\partial g}{\partial x_k}, \quad (3.1)$$

if we denote by $\omega^{j,k}$ the coefficients of ω^{-1} in local Darboux coordinates x_j .

Recall that for a holomorphic line bundle L with a fixed choice of Hermitian metric h , there is a unique connection, the “Chern connection”, which is compatible with both the metric and the holomorphic structure in a suitable sense [Huy, Prop. 4.2.14]. If we locally represent h by a matrix-valued function, the curvature of this connection is given by $\bar{\partial} \partial \log h$.

Definition 3.2 ([BeSl1, §2.1]). A compact Kähler manifold (M, ω) is **quantizable** if there is a holomorphic Hermitian line bundle (L, h) over M such that the curvature $\bar{\partial}\partial \log h$ of the Chern connection satisfies the **prequantum condition**

$$\bar{\partial}\partial \log h = -\sqrt{-1}\omega. \quad (3.2)$$

Then (L, h) gives a **quantization** of the Kähler manifold (M, ω) .

Condition (3.2) is there to ensure ensures the following.

Lemma 3.3 ([Schl2], [BeSl1, §2.1]). *Every quantization (L, h) of a compact Kähler manifold (M, ω) gives an embedding of M as a submanifold of \mathbb{CP}^{n-1} for some $n \in \mathbb{N}$, and hence M can be regarded as a projective algebraic variety (by Chow’s theorem). Conversely, every smooth projective algebraic variety is a quantizable compact Kähler manifold.*

Example 3.4 (The hyperplane bundle). The “tautological line bundle” over \mathbb{CP}^{n-1} is the holomorphic line bundle, usually denoted by $\mathcal{O}(-1)$, whose fiber over a point in \mathbb{CP}^{n-1} is the corresponding line in \mathbb{C}^n . The dual $\mathcal{O}(1) := \mathcal{O}(-1)^*$ of this line bundle is the **hyperplane line bundle** over \mathbb{CP}^{n-1} . The triple $(\mathbb{CP}^{n-1}, \mathcal{O}(1), \omega_{\text{FS}})$, where ω_{FS} is the Fubini–Study form and $\mathcal{O}(1)$ is equipped with the Fubini–Study metric, is the prototypical example of a quantizable Kähler manifold (M, L, ω) [Schl1].

The embedding into \mathbb{CP}^{n-1} mentioned in Lemma 3.3 requires a sufficiently positive line bundle, and (3.2) says only that L is positive (“ample”). It may therefore be necessary to use some tensor power $L^{\otimes m}$ of the line bundle L . However, by replacing L by $L^{\otimes m}$ and rescaling the Kähler form ω to $m\omega$ we can, and shall, assume that L is itself sufficiently positive (“very ample”).

The important requirement in Lemma 3.3 is that M *admits* a Kähler metric, but we do not need to choose one in order to an embedding of M into \mathbb{CP}^{n-1} as a complex-analytic submanifold (similarly we do not have to choose a Hermitian metric on L). Also, if we choose a Kähler metric ω on M and a Hermitian metric h on L , so that we can embed M as a Kähler submanifold of \mathbb{CP}^{n-1} , there is no need to require the prequantum relation (3.2) between h and ω ; it is just the existence of metrics satisfying the prequantum condition which is needed, to ensure that there is an ample line bundle on M . Therefore, we will often speak of a **polarized manifold**, i.e. a pair (M, L) where M is a compact Kähler manifold (with no choice of Kähler metric) and L is a positive line bundle on M (which we shall assume very ample for convenience, with no choice of Hermitian metric specified).

Let (M, ω) be a compact Kähler manifold and let (L, h) be a Hermitian line bundle over M . The space $H^0(M; L)$ of global holomorphic sections of L is finite-dimensional and hence made into a Hilbert space after fixing any inner product on $H^0(M; L)$. In Berezin quantization one looks at the limit of large m for the spaces $H^0(M; L^{\otimes m})$ of sections of the tensor powers of L , and therefore the inner products on these spaces should be comparable in some way. A consequence of the fact that the Kähler form ω is symplectic is that $\omega^d/d!$ (where $d := \dim_{\mathbb{C}} M$) is a volume form on M (the “Liouville form”) and we can take the inner product

$$\langle \phi | \psi \rangle_{h, \omega} := m^d \int_M h^m(\phi(x), \psi(x)) \frac{\omega(x)^d}{d!}, \quad \forall \psi, \phi \in H^0(M, L^m), \quad (3.3)$$

where h^m is the Hermitian metric on $L^{\otimes m}$ induced by h . Note that (3.3) is determined for all m by the choice of inner product on $H^0(M; L)$. If (L, h) is a quantization of (M, ω) then it is reasonable to leave out either h or ω from the notation in $\langle \cdot | \cdot \rangle_{h, \omega}$.

Example 3.5 (Sections of the hyperplane bundle). Recall the hyperplane line bundle $\mathcal{O}(1)$ over $\mathbb{P}[\mathfrak{H}^*]$ introduced in Example 3.4. Fixing a basis for $\mathfrak{H}^* \cong \mathbb{C}^n$, the global holomorphic sections of the m th tensor power $\mathcal{O}(m)$ of $\mathcal{O}(1)$ are identified with the degree- m homogeneous polynomials on \mathbb{C}^n . In particular, the space of holomorphic sections of $\mathcal{O}(1)$ is just the space \mathfrak{H} of continuous linear functionals on \mathfrak{H}^* . If we define \mathfrak{H}_m to be the Hilbert space of holomorphic sections of $\mathcal{O}(m)$ with inner product (3.3) then for $L = \mathcal{O}(1)$ we simply have $\mathfrak{H}_m = \mathfrak{H}^{\vee m}$ (symmetrized tensor product). Thus, the \mathfrak{H}_m ’s form the full symmetric subproduct system (see Example 2.2).

Let (L, h) be a quantization of (M, ω) and endow $H^0(M; L^m)$ with the inner product (3.3). The Hilbert space $L^2(M; L^{\otimes m})$ of all square-integrable sections of the line bundle $L^{\otimes m}$ is much larger than the holomorphic subspace $H^0(M; L^m)$. If Π_m denotes the projection from $L^2(M; L^{\otimes m})$ to $H^0(M; L^m)$ then every function $f \in C(M)$ defines an operator $\zeta^{(m)}(f)$ on $H^0(M; L^m)$ by

$$\zeta^{(m)}(f)\phi := \Pi_m(f\phi), \quad \forall \phi \in H^0(M; L^m), \quad (3.4)$$

i.e. acting as multiplication by f (recall that a section of a line bundle can be multiplied by continuous functions to yield a new section) followed by projection back to $H^0(M; L^m)$ (the latter step is needed since f is not holomorphic unless it is constant). In the case (L, h) is a quantization of (M, ω) , we have the following.

Proposition 3.6 ([BMS, Thms. 4.1, 4.2, §5]). *Let (L, h) be a quantization of a compact Kähler manifold (M, ω) and define a structure of Hilbert space on $H^0(M; L^m)$ by (3.3). Then the collection of endomorphism algebras $\text{End}_{\mathbb{C}} H^0(M; L^m)$ and maps $C^\infty(M) \ni f \rightarrow \zeta^{(m)}(f) \in \text{End}_{\mathbb{C}} H^0(M; L^m)$ defined by (3.4) gives a strict quantization of the algebra $C^\infty(M)$ in the sense of [Lan1, Def. 1.1.1], i.e. for all $f, g \in C^\infty(M)$ we have*

- (i) $\lim_{m \rightarrow \infty} \|\zeta^{(m)}(f)\| = \|f\|$ (Rieffel condition),
- (ii) $\lim_{m \rightarrow \infty} \|\zeta^{(m)}(fg) - \zeta^{(m)}(f)\zeta^{(m)}(g)\| = 0$ (von Neumann condition),
- (iii) $\lim_{m \rightarrow \infty} \|m^{-1}[\zeta^{(m)}(f), \zeta^{(m)}(g)] - \{f, g\}\| = 0$ (Dirac condition),

and every operator in $\text{End}_{\mathbb{C}} H^0(M; L^m)$ is of the form $\zeta^{(m)}(f)$ for some $f \in C^\infty(M)$ [BMS, Prop. 4.2]. Here $\{\cdot, \cdot\}$ is the Poisson bracket (3.1).

For any volume form $\omega^d/d!$ of (M, L) , we have the associated Lebesgue space $L^2(M, \omega)$. The algebra $C(M)$ identifies with a subspace of $L^2(M, \omega)$. Let $\varsigma^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow C(M)$ denote the adjoint of $\zeta^{(m)} : C(M) \rightarrow \mathcal{B}(\mathfrak{H}_m)$ with respect to the L^2 -inner product on $L^2(M, \omega)$ and the normalized Hilbert-Schmidt inner product on $\mathcal{B}(\mathfrak{H}_m)$.

For $A \in \mathcal{B}(\mathfrak{H}_m)$, the function $\varsigma^{(m)}(A)$ is also called the **Berezin covariant symbol** of A , and if $A = \zeta^{(m)}(f)$ then f is a (non-unique) **contravariant symbol** of A . The map

$$\varsigma^{(m)} \circ \zeta^{(m)} : C^\infty(M) \rightarrow C^\infty(M)$$

is the **Berezin transform** at level m .

Since the Toeplitz maps $\zeta^{(m)}$ are surjective by Proposition 3.6, their adjoints $\varsigma^{(m)}$ are injective. Hence we may regard the $\mathcal{B}(\mathfrak{H}_m)$'s as embedded in $C^\infty(M)$. One may then ask if it is possible to use the covariant symbols to approximate $C^\infty(M)$ by finite-dimensional matrix algebras $\mathcal{B}(\mathfrak{H}_m)$.

Similar to the famous expansion of the integral kernel for the Bergman projections Π_m [Zeld1], the Berezin transform $\varsigma^{(m)} \circ \zeta^{(m)}$ has an asymptotic expansion at large m [Sch1]. For homogeneous polarized manifolds (M, L) (namely coadjoint orbits for compact Lie groups), the Berezin transforms converge to the identity map on $C(M)$ as m goes to infinity [Rie2, Thm. 6.1]. This convergence result, which is stronger than the mere Toeplitz convergence in Proposition 3.6, relies on the fact coadjoint orbits are “balanced” (see below).

We shall see that by changing the definition of the Toeplitz maps $\zeta^{(m)}$ we can in fact obtain $C^\infty(M)$ as an inductive limit for *any* polarized manifold (M, L) .

3.2 Projectively induced quantization

The vector spaces $H^0(M; L^m)$ equipped with the inner products (3.3) do not always form a sub-product system. For that one has to choose ω and h appropriately, and for most polarized manifolds (M, L) one cannot choose them to satisfy the prequantum condition (3.2) at the same time.

If we do *not* require that h and ω are related as in (3.2) then any two of (i) an inner product on $H^0(M; L)$, (ii) a Hermitian metric on L and (iii) a volume form $\omega^d/d!$ on M determines the third via (3.3). By the Calabi–Yau theorem [Yau1], any volume form on M can be obtained as $\omega^d/d!$ for

some ω in the cohomology class $c_1(L)$ (for any choice of polarization L , after normalization of the volume form).

For us, the choice of inner product $\langle \cdot | \cdot \rangle$ on $H^0(M; L)$ will be the important input, and it will not matter which Hermitian metric on L and volume form on M was used to define it.

Given a polarized manifold (M, L) , a choice of basis for the n -dimensional vector space $H^0(M; L)$ allows use to embed M into $\mathbb{P}[H^0(M; L)^*]$. The elements of the basis for $H^0(M; L)$ become the restrictions of the homogeneous coordinate functions z_1, \dots, z_n on $\mathbb{P}[H^0(M; L)^*]$ to the embedded M . Choosing an inner product $\langle \cdot | \cdot \rangle$ on $H^0(M; L)$ we obtain an n -dimensional Hilbert space \mathfrak{H} which after a choice of orthonormal basis identifies with \mathbb{C}^n , and so M embeds into $\mathbb{P}[\mathfrak{H}^*] = \mathbb{CP}^{n-1}$. Whatever inner product on $H^0(M; L)$ we used to define the Hilbert space \mathfrak{H} , it will produce the symmetric subproduct system $\mathfrak{H}^{\vee \bullet}$ of holomorphic sections of the hyperplane bundle on $\mathbb{P}[\mathfrak{H}^*]$ as in Example 3.5. What Lemma 3.3 says is that the ideal determined by the algebraic relations among the z_j 's, appearing when we restrict them to the submanifold M , is homogeneous. The subspaces $H^0(M; L^m) \subset H^0(M; L)^{\vee m}$ of holomorphic sections of the tensor powers of L endowed with the inner product as a subspace of $\mathfrak{H}^{\vee m}$ will be denoted by

$$\mathfrak{H}_m = (H^0(M; L), \langle \cdot | \cdot \rangle).$$

Here $\langle \cdot | \cdot \rangle$ is thus the inner product (3.3) in the special case when ω and h are the restrictions to M of the Fubini–Study metrics on $\mathbb{P}[\mathfrak{H}^*]$, depending only on the inner product on $H^0(M; L)$ which defines the one-particle Hilbert space \mathfrak{H} . We set $\mathfrak{H}_0 := \mathbb{C}$.

We therefore have a description of polarized manifolds (M, L) with the extra datum of an inner product on $H^0(M; L)$ as a collection of Hilbert spaces \mathfrak{H}_m satisfying (2.1) with $\mathfrak{H}_m \subseteq \mathfrak{H}^{\vee m}$, where \vee is the symmetrized tensor product (recall Lemma 2.9). The subproduct system \mathfrak{H}_\bullet is obtained from $\mathfrak{H}^{\vee \bullet}$ by quotienting out by the ideal in $\mathbb{C}[z_1, \dots, z_n]$ which defines the embedded M .

Corollary 3.7. *Every commutative subproduct system $\mathfrak{H}_\bullet = (\mathfrak{H}_m)_{m \in \mathbb{N}_0}$ (see Definition 2.3) determines (via its associated homogeneous ideal in $\mathbb{C}[z_1, \dots, z_n]$) a polarized manifold (M, L) with a fixed structure of Hilbert space on $H^0(M; L)$. Conversely, every such datum $(M, L, \langle \cdot | \cdot \rangle)$ determines a commutative subproduct system.*

We stress that for obtaining the subproduct system \mathfrak{H}_\bullet , the inner product on $H^0(M; L)$ is arbitrary; we do not require h and ω in (3.3) to satisfy the pre-quantum condition. Even if we did require $\omega = \sqrt{-1} \bar{\partial} \partial \log h$, the inner product on $\mathfrak{H}_m \subset \mathfrak{H}^{\vee m}$ for $m \geq 2$ would in general differ from the inner product $\langle \cdot | \cdot \rangle_h$ defined by the initial ω and h .

Lemma 3.8. *Let (M, L) be a polarized manifold with $d := \dim_{\mathbb{C}} M$. Then for any inner product $\langle \cdot | \cdot \rangle$ on $H^0(M; L)$, there exists a unique volume form on M , which we can express as ω^d for a Kähler metric in the class $c_1(L)$, such that $\langle \cdot | \cdot \rangle$ is ω^d -balanced in the sense of [Don3, §2.2], i.e. if Z_1, \dots, Z_n are the homogeneous coordinates on $M \subset \mathbb{P}[\mathfrak{H}^*]$ associated with any orthonormal basis for $\mathfrak{H} = (H^0(M; L), \langle \cdot | \cdot \rangle)$, normalized to $\sum_{k=1}^n Z_k Z_k^* = \mathbf{1} \in C^\infty(M)$, then*

$$\frac{1}{\text{vol}(M, L)} \int_M Z_j^* Z_k \frac{\omega^d}{d!} = \frac{\delta_{j,k}}{n}, \quad \forall j, k = 1, \dots, n,$$

where $\text{vol}(M, L) := \int_M \omega^d / d!$. In that case one also has (using Notation 2.12) for all $\mathbf{j}, \mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{j}| = m = |\mathbf{k}|$ that

$$\frac{1}{\text{vol}(M, L)} \int_M Z_{\mathbf{j}}^* Z_{\mathbf{k}} \frac{\omega^d}{d!} = \frac{\delta_{\mathbf{j}, \mathbf{k}}}{\text{Tr}(p_m)},$$

where $p_m : \mathfrak{H}^{\otimes m} \rightarrow \mathfrak{H}_m$ is the orthogonal projection and $\mathfrak{H}_\bullet \subset \mathfrak{H}^{\vee \bullet}$ is the subproduct system of $(M, L, \langle \cdot | \cdot \rangle)$.

Proof. The first part follows from the Calabi–Yau theorem [Yau1] and the fact that every polarized manifold (M, L) admits a unique ω^d -balanced metric for every volume form ω^d on M [BLY1], [Don3, §2.2] (equivalently, every line bundle is stable and from this it follows that every very ample line bundle is “balanced as a line bundle” in the sense of [Wa1]). The second statement is immediate from the definition of \mathfrak{H}_\bullet . \square

To say that (M, L) is **balanced** as a polarized manifold [Don1] means precisely that there exists a Hilbert space structure \mathfrak{H} on $H^0(M; L)$ such that the volume form ω^d in Lemma 3.8 is the restriction to M of the Fubini–Study volume form on $\mathbb{P}[\mathfrak{H}^*]$. Not every polarized manifold is balanced, and so in general one cannot form a subproduct system from a quantization (L, h) of a compact Kähler manifold (M, ω) in the sense of the last section. We consider therefore instead the following quantization.

Definition 3.9. A **projectively induced** quantization of a polarized manifold (M, L) is the datum of a subproduct system $\mathfrak{H}_\bullet \subseteq \mathfrak{H}^{\vee\bullet}$ associated with some choice of inner product $\langle \cdot | \cdot \rangle$ on $H^0(M; L)$, together with the covariant and contravariant symbol maps specified by \mathfrak{H}_\bullet . That is, we regard $C(M)$ as a subspace of the Hilbert space $L^2(M; \omega)$ of square-integrable functions on M with respect to the $\langle \cdot | \cdot \rangle$ -balancing volume form guaranteed by Lemma 3.8, and we take the Toeplitz map $\zeta^{(m)} : C^\infty(M) \rightarrow \mathcal{B}(\mathfrak{H}_m)$ defined by

$$\zeta^{(m)}(f)\phi := \Pi_m(f\phi), \quad \forall \phi \in \mathfrak{H}_m,$$

where $\Pi_m : L^2(M; L^m) \rightarrow \mathfrak{H}_m$ is the orthogonal projection, and we let $\zeta^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow C(M) \subset L^2(M; \omega)$ be its adjoint. The Hermitian metric h on L used to define $L^2(M; L^m)$ is then forced to be the Fubini–Study metric associated with $\langle \cdot | \cdot \rangle$ (cf. [Don2]), i.e. the unique Hermitian metric $h = \text{FS}(\langle \cdot | \cdot \rangle)$ on L for which any orthonormal basis Z_1, \dots, Z_n for \mathfrak{H} satisfies

$$\sum_{k=1}^n h(Z_k, Z_k) = \mathbf{1} \in C(M).$$

An explicit formula for the covariant symbol map $\zeta^{(m)}$ is easy to write down; see Theorem 4.15.

Remark 3.10. The terminology in Definition 3.9 is slightly nonstandard unless (M, L) is balanced; indeed, (M, L) is a balanced polarized manifold (in the sense of [Don1]) if and only if there exists a Hermitian metric h on L such that the quantization (L, h) of $(M, \bar{\partial}\partial \log h)$ is projectively induced for some choice of inner product on $H^0(M; L)$. In that sense the term “projectively induced” appeared in [CGR1] (it says precisely that the epsilon function discussed there is constant; a less illuminating name for the same thing is a “regular” quantization [CGR2]). In the less restricted sense of Definition 3.9, which works for any polarized manifold (M, L) since we do not require the prequantization condition, the projectively induced quantizations were referred to as “Berezin–Bergman quantizations” in [LMS1, §5] (this is the only work we know of where it has been discussed for not necessarily balanced manifolds).

We shall see that using projectively induced quantizations, a polarized manifold (M, L) need not be balanced in order to recover $C^\infty(M)$ using Berezin quantization. We stress that the restriction to quantizations with this choice of volume form is not an artifact of the operator-algebraic approach: it is needed if we want the stronger convergence of Berezin transforms, and in order to recover the algebra $C^\infty(M)$ not just approximately.

3.3 The circle bundle

Recall the Toeplitz quantization maps $\zeta^{(m)} : C(M) \rightarrow \mathcal{B}(\mathfrak{H}_m)$ (for definiteness and later relevance we will focus on the case of a projectively induced quantization). We can assemble them into a single map

$$\zeta : C(M) \rightarrow \prod_{m \in \mathbb{N}_0} \mathcal{B}(\mathfrak{H}_m), \quad (3.5)$$

where the C^* -algebra on the right-hand side is the C^* -direct product.

Let L^* be the dual line bundle of L . Under the embedding of M into \mathbb{CP}^{n-1} , when L becomes the restriction of the hyperplane line bundle, L^* becomes the restriction of the tautological line bundle. Denote by

$$\mathbb{S}_M := \{\zeta \in L^* \mid \|\zeta\| = 1\} \quad (3.6)$$

the total space of the associated principal $U(1)$ -bundle. The $U(1)$ -action on \mathbb{S}_M induces a \mathbb{Z} -grading

$$C(\mathbb{S}_M) = \overline{\bigoplus_{k \in \mathbb{Z}} C(\mathbb{S}_M)^{(k)}}^{\|\cdot\|}$$

of the C^* -algebra of continuous functions on \mathbb{S}_M , with $C(\mathbb{S}_M)^{(0)} = C(M)$. The C^* -algebra $C(\mathbb{S}_M)$ is generated by the normalized homogeneous coordinate functions Z_1, \dots, Z_n on $M \subset \mathbb{CP}^{n-1}$. Products of the form $Z_{j_1} \cdots Z_{j_m} Z_{k_1}^* \cdots Z_{k_l}^*$ constitute a basis for the L^2 -space $L^2(\mathbb{S}_M)$ (see [Schl1, §5.1] for the definition of the measure on \mathbb{S}_M) and we define the “Hardy space” $H^2(\mathbb{S}_M)$ to be the subspace of $L^2(\mathbb{S}_M)$ spanned by the products $Z_{j_1} \cdots Z_{j_m}$ for all $m \in \mathbb{N}_0$. The Hardy space can therefore be obtained as the Hilbert space direct sum

$$H^2(\mathbb{S}_M) = \overline{\bigoplus_{m \in \mathbb{N}_0} \mathfrak{H}_m}^{\langle \cdot | \cdot \rangle_{L^2}},$$

just as the Fock space $\mathfrak{H}_{\mathbb{N}}$ associated with the subproduct system \mathfrak{H}_{\bullet} . The inner products differ slightly, the inner product of two elements $\psi, \varphi \in \mathfrak{H}_m \subset \mathfrak{H}_{\mathbb{N}}$ being related to their inner product as elements of $H^2(\mathbb{S}_M)$ as

$$\langle \psi | \varphi \rangle_{L^2} = \frac{1}{\text{Tr}(p_m)} \langle \psi | \varphi \rangle,$$

where $p_m \in \mathcal{B}(\mathfrak{H}_m)$ is the identity operator. So Fock space can be embedded into $H^2(\mathbb{S}_M)$,

$$\mathfrak{H}_{\mathbb{N}} = \overline{\bigoplus_{m \in \mathbb{N}_0} \mathfrak{H}_m}^{\langle \cdot | \cdot \rangle} \subset H^2(\mathbb{S}_M) \subset L^2(\mathbb{S}_M).$$

Let $\Pi : L^2(\mathbb{S}_M) \rightarrow \mathfrak{H}_{\mathbb{N}}$ be the orthogonal projection. Then, with ζ as in (3.5), we have

$$\zeta(f)\psi = \Pi(f\psi), \quad \forall \psi \in \mathfrak{H}_{\mathbb{N}},$$

if we identify $f \in C(M)$ with the multiplication operator it defines on $L^2(\mathbb{S}_M)$. Then f is just the **contravariant symbol** of the operator $\zeta(f)$ in the general sense of [Bere3]. The interior of $\mathbb{S}_{\mathbb{M}}$ (the disk bundle) is a bounded symmetric domain and Berezin quantization on spaces such as \mathbb{S}_M has been studied even more than in the setting of compact Kähler manifolds, see e.g. [UnUp1], [Bere2].

More details about the circle bundle can be found in [Schl1, §2], [Zeld1, §2].

We can define the Berezin transform and the covariant symbol of an operator on $\mathfrak{H}_{\mathbb{N}}$, just as we did on the components \mathfrak{H}_m , using the fact that $\mathfrak{H}_{\mathbb{N}}$ is a reproducing-kernel Hilbert space. Again the covariant symbol map ς is the adjoint of ζ . In the noncommutative setting we will just calculate the adjoint of ζ and take that as the definition of the covariant symbol map.

3.4 Singular varieties

We have seen that Berezin quantization of quantizable Kähler manifolds is really the quantization of smooth projective varieties. It was suggested in [Schl2] that it may be possible to quantize also singular (non-smooth) projective varieties in the same fashion. We shall see that this is in fact so: it will be covered by the constructions in the next two sections, as the case when the subproduct system \mathfrak{H}_{\bullet} is commutative (Corollary 5.31).

4 Inductive limits

Recall [AraM1, Def. 1.1.11] that a sequence $(\mathcal{B}_m)_{m \in \mathbb{N}_0}$ of $*$ -algebras \mathcal{B}_m forms an “inductive system” if there are $*$ -homomorphisms $\iota_{m,l} : \mathcal{B}_m \rightarrow \mathcal{B}_{m+l}$ for each $m \leq l$, and that the **algebraic inductive**

limit of such a sequence is the algebra $\bigcup_{m \in \mathbb{N}_0} \mathcal{B}_m$ obtained as the quotient of the algebra of eventually constant sequences of elements in the \mathcal{B}_m 's,

$$\left\{ (b_j)_{j \in \mathbb{N}_0} \in \prod_{j \in \mathbb{N}_0} \mathcal{B}_j \mid \exists m \in \mathbb{N}_0 \text{ such that } b_j = b_m \text{ for all } j \geq m \right\},$$

by its ideal of sequences $(b_j)_{j \in \mathbb{N}_0}$ which are eventually 0. If each \mathcal{B}_m is a C^* -algebra, $\bigcup_{m \in \mathbb{N}_0} \mathcal{B}_m$ can be completed in a canonical C^* -norm to obtain a C^* -algebra which, if the $\iota_{m,l}$'s are injective, can be identified with the non-disjoint union $\overline{\bigcup_{m \in \mathbb{N}_0} \mathcal{B}_m}^{\|\cdot\|}$ [AraM1, §1.2].

It was observed in [Hawk1] that the sequence of algebras $\mathcal{B}_m := \mathcal{B}(\mathfrak{H}_m)$ arising in quantization has a structure resembling that of an inductive system, although the map from \mathcal{B}_m to \mathcal{B}_{m+1} is not a homomorphism in the category of C^* -algebras. If we want to obtain a C^* -algebra $C(M)$ of continuous functions on a manifold as an inductive limit of finite-dimensional matrix algebras, then requiring $\mathcal{B}_m \subset \mathcal{B}_{m+1}$ says by definition that $C(M)$ is an AF algebra. This forces M to be totally disconnected. Hence we must relax the notion of inductive limit.

4.1 Relaxed definition of inductive limits

Blackadar and Kirchberg introduced a more general inductive-limit-type construction [BKil]. Although never pointed out in the literature, the system of finite-dimensional C^* -algebras $\mathcal{B}(\mathfrak{H}_m)$ obtained from a projective quantization (M, ω, L) fits perfectly into their framework. This is most apparent in [Hawk1] where similar notions were introduced independently. We will follow the notation of [Hawk1] as closely as possible.

Notation 4.1. If $\mathcal{B}_\bullet = (\mathcal{B}_m)_{m \in \mathbb{N}_0}$ is a sequence of C^* -algebras, we write

$$\Gamma_b(\mathcal{B}_\bullet) = \prod_{m \in \mathbb{N}_0} \mathcal{B}_m$$

for the full C^* -**direct product** of the \mathcal{B}_m 's, i.e. the set of sequences $X_\bullet = (X_m)_{m \in \mathbb{N}_0}$ of elements $X_m \in \mathcal{B}_m$ with finite supremum norm

$$\|X_\bullet\| := \sup_{m \in \mathbb{N}_0} \|X_m\|_{\mathcal{B}_m} < \infty.$$

The multiplication and $*$ -operation in $\Gamma_b(\mathcal{B}_m)$ is pointwise. We also write

$$\Gamma_0(\mathcal{B}_\bullet) = \overline{\bigoplus_{m \in \mathbb{N}_0} \mathcal{B}_m}$$

for the C^* -**direct sum**, the closed two-sided ideal in $\Gamma_b(\mathcal{B}_\bullet)$ consisting of the sequences converging to zero in norm. We simply write $\Gamma_b := \Gamma_b(\mathcal{B}_\bullet)$ etc. if it is clear which sequence \mathcal{B}_\bullet it concerns. We let

$$\pi : \Gamma_b \rightarrow \Gamma_b / \Gamma_0$$

be the quotient map

Since we will only deal with a special kind of the “generalized inductive systems” defined in [BKil] (namely the “NF” ones), we will simply refer to them as “inductive systems”. See also [BrOz1, §11].

Definition 4.2. An **inductive system** is a sequence $(\mathcal{B}_\bullet, \iota_\bullet)$ of full matrix algebras $\mathcal{B}_m = M_{k(m)}(\mathbb{C})$ and unital completely positive maps $\iota_{m,l} : \mathcal{B}_m \rightarrow \mathcal{B}_{m+l}$ for $l \geq m$ (with $\iota_{m,m} := \text{id}$) satisfying

$$\iota_{m,l} = \iota_{r,l} \circ \iota_{m,r}, \quad \text{if } m \leq r \leq l \tag{4.1}$$

and which are **asymptotically multiplicative** in the sense that for all $A, B \in \mathcal{B}(\mathfrak{H}_m)$, $\varepsilon > 0$, there are $r \leq l$ such that

$$\|\iota_{r,l}(\iota_{m,r}(A)\iota_{m,r}(B)) - \iota_{m,l}(A)\iota_{m,l}(B)\| < \varepsilon. \tag{4.2}$$

The **inductive limit** of an inductive system $(\mathcal{B}_\bullet, \iota_\bullet)$ is the C^* -algebra

$$\mathcal{B}_\infty \subset \Gamma_b(\mathcal{B}_\bullet)/\Gamma_0(\mathcal{B}_\bullet)$$

generated by the elements

$$\varsigma^{(m)}(A) := \pi((\iota_l(A))_{l \geq m}), \quad A \in \mathcal{B}(\mathfrak{H}_m) \quad (4.3)$$

for all $m \in \mathbb{N}_0$, where $\pi : \Gamma_b \rightarrow \Gamma_b/\Gamma_0$ is the quotient map.

Remark 4.3 (Norm). A norm on the quotient C^* -algebra Γ_b/Γ_0 is given by

$$\|\pi(X_\bullet)\| = \limsup_{m \rightarrow \infty} \|X_m\|, \quad \forall X_\bullet \in \Gamma_b,$$

and this norm satisfies the C^* -identity, hence it is the unique C^* -norm on Γ_b/Γ_0 . Moreover, since the $\iota_{m,l}$'s are norm-decreasing,

$$\limsup_{m \rightarrow \infty} \|\varsigma^{(m)}(A)\| = \lim_{l \rightarrow \infty} \|\iota_{m,l}(A)\|, \quad \forall A \in \mathcal{B}(\mathfrak{H}_m),$$

so the norm on \mathcal{B}_∞ is just the “norm-at-infinity” of $\pi^{-1}(\mathcal{B}_\infty)$.

It follows that the maps

$$\varsigma^{(m)} : \mathcal{B}_m \rightarrow \mathcal{B}_\infty \quad (4.4)$$

are completely positive, and we refer to $\varsigma^{(m)}$ as the **covariant Berezin symbol map** at level m . The motivation for this terminology will become clear below. Due to (4.1), the covariant symbol maps satisfy

$$\varsigma^{(l)} \circ \iota_{m,l} = \varsigma^{(m)}, \quad \forall m \leq l \in \mathbb{N}_0. \quad (4.5)$$

We may say that a sequence $A_\bullet \in \Gamma_b$ is **eventually constant** under ι_\bullet if there is a large enough $m \in \mathbb{N}_0$ such that $A_l = \iota_{r,l}(A_r)$ for all $l \geq r \geq m$. Then \mathcal{B}_∞ is the image under π of the norm closure of the algebra of eventually constant sequences.

Remark 4.4 (Asymptotic multiplicativity). The condition (4.2) is chosen precisely to ensure that $\varsigma^{(m)}(A)\varsigma^{(m)}(B)$ belongs to the C^* -algebra \mathcal{B}_∞ for all $A, B \in \mathcal{B}(\mathfrak{H}_m)$, without requiring it to be close to $\varsigma^{(m)}(AB)$. Conversely, if $\varsigma^{(m)}(A)\varsigma^{(m)}(B)$ belongs to \mathcal{B}_∞ for all A and B then each $\varsigma^{(m)}(A)\varsigma^{(m)}(B)$ is an eventually constant sequence, so (4.2) must hold.

Remark 4.5 (Continuous fields). Suppose that $\mathcal{B}(\bullet)$ is a continuous field of matrix algebras over $\mathbb{N}_0 \cup \{\infty\}$. For each $A \in \mathcal{B}(\infty)$, there is a continuous section $x \rightarrow A(x)$ of $\mathcal{B}(\bullet)$ with $A(\infty) = A$, and this section defines an element of $\prod_{m \in \mathbb{N}_0} \mathcal{B}(m)$. Two sections evaluating to A at ∞ differ by an element in $\bigoplus_{m \in \mathbb{N}_0} \mathcal{B}(m)$. Hence [BlKi1, Prop. 2.2.3]

$$\mathcal{B}(\infty) \subset \left(\prod_{m \in \mathbb{N}_0} \mathcal{B}(m) \right) / \left(\bigoplus_{m \in \mathbb{N}_0} \mathcal{B}(m) \right).$$

In fact, a C^* -algebra is an inductive limit (in the sense of Definition 4.2) if and only if it is a nuclear separable C^* -algebra which is of the form $\mathcal{B}(\infty)$ for some continuous field of matrix algebras over $\mathbb{N}_0 \cup \{\infty\}$ [BlKi1, Thm. 5.2.2].

Remark 4.6 (Quasi-diagonality). If a C^* -algebra \mathcal{B}_∞ is an inductive limit in the sense of Definition 4.2, there exists a short-exact sequence

$$0 \longrightarrow \mathcal{K} \longrightarrow \mathcal{D} \longrightarrow \mathcal{B}_\infty$$

which is an “essential quasi-diagonal extension” of \mathcal{B}_∞ , meaning that \mathcal{D} is a C^* -algebra of quasi-diagonal operators containing the C^* -algebra \mathcal{K} of compact operators as an essential ideal. In fact, such an extension of a C^* -algebra \mathcal{B}_∞ exists if and only if \mathcal{B}_∞ can be embedded into $\Gamma_b(\mathcal{B}_\bullet)/\Gamma_0(\mathcal{B}_\bullet)$ for some sequence of full matrix algebras \mathcal{B}_m [BlKi1].

Remark 4.7 (Nuclearity). Since \mathcal{B}_∞ is nuclear, the Choi–Effros lifting theorem [Blac1, IV.2.3.4] says that the identity mapping $\text{id} : \mathcal{B}_\infty \rightarrow \mathcal{B}_\infty$ can be lifted to a unital completely positive map $\zeta : \mathcal{B}_\infty \rightarrow \Gamma_b$ such that if

$$\zeta^{(m)}(f) := \zeta(f)p_m \in \mathcal{B}(\mathfrak{H}_m), \quad \forall f \in \mathcal{B}_\infty,$$

then the sequence $\zeta^{(m)} \circ \zeta$ converges in the point-norm topology to $\text{id} : \mathcal{B}_\infty \rightarrow \mathcal{B}_\infty$. We shall calculate ζ and its inverse ς explicitly in §5.6.

4.2 Inductive limits from subproduct systems

Theorem 4.8. *Every subproduct system \mathfrak{H}_\bullet defines a generalized inductive system $(\mathcal{B}_\bullet, \iota_{\bullet,\bullet})$ by setting $\mathcal{B}_m := \mathcal{B}(\mathfrak{H}_m)$ and*

$$\iota_{m,l}(A) := p_l(A \otimes \mathbf{1}_{\mathfrak{H} \otimes (l-m)})p_l, \quad \forall A \in \mathcal{B}(\mathfrak{H}_m), \quad (4.6)$$

where $p_l : \mathfrak{H}^{\otimes l} \rightarrow \mathfrak{H}_l$ is the projection.

Proof. It is clear that each $\iota_{m,l} : \mathcal{B}_m \rightarrow \mathcal{B}_l$ is unital and completely positive. For $m \leq r \leq l$ and $A \in \mathcal{B}(\mathfrak{H}_m)$ we have

$$\begin{aligned} \iota_{r,l} \circ \iota_{m,r}(A) &= \iota_{m,l}(p_r(A \otimes \mathbf{1}_{\mathfrak{H} \otimes (r-m)})p_r) = p_l(p_r(A \otimes \mathbf{1}_{\mathfrak{H} \otimes (r-m)})p_r \otimes \mathbf{1}_{\mathfrak{H} \otimes (l-r)})p_l \\ &= p_l(A \otimes \mathbf{1}_{\mathfrak{H} \otimes (l-m)})p_l, \end{aligned}$$

where the last equality is due to (2.2). That is, the coherence condition (4.1) holds.

It remains to show that $\iota_{\bullet,\bullet}$ is asymptotically multiplicative, i.e. that it satisfies (4.2). We have

$$\begin{aligned} \iota_{r,l}(\iota_{m,r}(A)\iota_{m,r}(B)) &= \iota_{r,l}(p_r(A \otimes \mathbf{1})p_r(B \otimes \mathbf{1})p_r) \\ &= p_l((p_r(A \otimes \mathbf{1})p_r(B \otimes \mathbf{1})p_r) \otimes \mathbf{1})p_l \\ &= p_l(((A \otimes \mathbf{1})p_r(B \otimes \mathbf{1})) \otimes \mathbf{1})p_l, \end{aligned}$$

so it is the failure of $A \otimes \mathbf{1}$ to commute with the projection p_r which spoils multiplicativity. It seems hard to show directly from norm estimates that the maps (4.6) satisfy the asymptotic multiplicativity condition (4.2). We shall instead obtain that by showing that the set of elements (4.3) forms an algebra.

Consider the strongly graded ring

$$\mathcal{R} = \bigoplus_{m \in \mathbb{N}_0} \mathcal{R}_m := \bigoplus_{m \in \mathbb{N}_0} \mathcal{O}_{\mathfrak{H}}^{(m)}.$$

Denote by $\text{Gr}(\mathcal{R})$ the Abelian category of \mathbb{Z} -graded right \mathcal{R} -modules, with morphisms the grading-preserving morphisms in the category of \mathbb{R} -modules. Write $\mathcal{R}_{\geq m}$ for the \mathcal{R} -module $\bigoplus_{l \geq m} \mathcal{R}_l$. Then it is easy to see that

$$\text{End}_{\text{Gr}(\mathcal{R})}(\mathcal{R}_{\geq m}) = \mathcal{B}(\mathfrak{H}_m)$$

as rings, and that $\iota_{m,l}$ maps an element of $\text{End}_{\text{Gr}(\mathcal{R})}(\mathcal{R}_{\geq m})$ to its restriction to the submodule $\mathcal{R}_{\geq l} \subset \mathcal{R}_{\geq m}$. By [Sten1, §IX.1], the algebraic inductive limit

$${}^0\mathcal{B}_\infty = \lim_{m \rightarrow \infty} \text{End}_{\text{Gr}(\mathcal{R})}(\mathcal{R}_{\geq m})$$

is a ring (and an algebra since the maps $\iota_{m,l}$ are linear). Therefore the norm closure \mathcal{B}_∞ of ${}^0\mathcal{B}_\infty$ is an algebra as well. In particular, the asymptotic multiplicativity condition (4.2) is satisfied. \square

Thus, subproduct systems give rise to generalized inductive limits of C^* -algebras with the special property that the algebraic direct limit ${}^0\mathcal{B}_\infty$ is already an algebra (no need for norm closure). Still, the asymptotic multiplicativity condition (4.2) cannot be formulated in a weaker fashion even for subproduct systems, since the set of eventually constant sequences under $\iota_{\bullet,\bullet}$ is only an algebra after taking norm closure.

Our aim is to identify the inductive limit \mathcal{B}_∞ with the Cuntz–Pimsner core $\mathcal{O}_{\mathfrak{H}}^{(0)}$. For that we need two lemmas.

Lemma 4.9. *The completely positive maps $\iota_{m,l} : \mathcal{B}_m \rightarrow \mathcal{B}_l$ defined in (4.6) can be expressed as*

$$\iota_{m,l}(A) = \sum_{|\mathbf{k}|=l-m} R_{\mathbf{k}} A R_{\mathbf{k}}^*|_{\mathfrak{H}_l} \quad (4.7)$$

$$= \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_l} \quad (4.8)$$

for all $A \in \mathcal{B}(\mathfrak{H}_m)$, where $R_{\mathbf{k}}$ is the right shift by the vector $e_{\mathbf{k}}$ as in (2.6) and $A_{\mathbf{j},\mathbf{k}} := \langle e_{\mathbf{j}} | A e_{\mathbf{k}} \rangle$.

Proof. Formula (4.8) is immediate from

$$S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_l} = p_l(|p_m e_{\mathbf{j}}\rangle \langle p_m e_{\mathbf{k}}| \otimes \mathbf{1}_{\mathfrak{H}_{l-m}}) p_l.$$

Similarly, the expression (4.7) is deduced from straightforward calculations. \square

Lemma 4.10. *Let $\pi^{-1}(\mathcal{B}_{\infty})$ be the norm closure of the subset of $\prod_m \mathcal{B}(\mathfrak{H}_m)$ consisting of sequences which are eventually constant under the coherent system $\iota_{\bullet,\bullet}$ defined by (4.6). Then $\pi^{-1}(\mathcal{B}_{\infty})$ coincides with the normally ordered part of the Toeplitz core $\mathcal{T}_{\mathfrak{H}}^{(0)}$. Hence the normally ordered part of $\mathcal{T}_{\mathfrak{H}}^{(0)}$ is an algebra, and must coincide with all of $\mathcal{T}_{\mathfrak{H}}^{(0)}$, so*

$$\pi^{-1}(\mathcal{B}_{\infty}) = \mathcal{T}_{\mathfrak{H}}^{(0)}.$$

In this way we have proven Lemma 2.15.

Proof. It is clear that every normally ordered element of $\mathcal{T}_{\mathfrak{H}}^{(0)}$ defines a sequence $A_{\bullet} = (A_m)_{m \in \mathbb{N}_0}$ of operators $A_m \in \mathcal{B}(\mathfrak{H}_m)$ with $\iota_{m,l}(A_m) = A_l$ for sufficiently large $m \leq l$. For example,

$$S_{\mathbf{j}} S_{\mathbf{k}}^* = (S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_m})_{m \in \mathbb{N}_0} \in \prod_{m \in \mathbb{N}_0} \mathcal{B}(\mathfrak{H}_m).$$

Suppose now that $A = (A_m)_{m \in \mathbb{N}_0}$ is any element of $\prod_{m \in \mathbb{N}_0} \mathcal{B}(\mathfrak{H}_m)$ which is eventually constant. Since $\mathcal{B}(\mathfrak{H}_m)$ is contained in $\mathcal{T}_{\mathfrak{H}}^{(0)}$ for each m , we may for simplicity just as well look at the case where $\iota_{r,l}(A_r) = A_l$ for all $r \leq l$ for some $r \in \mathbb{N}_0$ while $A_m = 0$ for $m \leq r$. Then (4.8) in Lemma 4.9 shows that A is a combination of shift operators.

From the fact that \mathcal{B}_{∞} is an algebra we have that $\pi^{-1}(\mathcal{B}_{\infty})$ is an algebra, whence the last statement. \square

Remark 4.11. Let $\mathcal{B}(\bullet)$ be the continuous field of C^* -algebras over $\mathbb{N}_0 \cup \{\infty\}$ such that the fiber over $m \in \mathbb{N}$ is $\mathcal{B}(m) = \mathcal{B}(\mathfrak{H}_m)$ and the fiber over ∞ is $\mathcal{B}(\infty) = \mathcal{B}_{\infty}$, the inductive limit (cf. Remark 4.5). Then Lemma 4.10 says that $\mathcal{T}_{\mathfrak{H}}^{(0)}$ is the algebra of continuous sections of this field. For commutative case see also [Haw2, Thm. 3.3].

Theorem 4.12. *Let \mathfrak{H}_{\bullet} be a subproduct system and let $\mathcal{O}_{\mathfrak{H}}^{(0)}$ denote the $U(1)$ -invariant part of the Cuntz-Pimsner algebra of \mathfrak{H}_{\bullet} . Then we have*

$$\mathcal{O}_{\mathfrak{H}}^{(0)} \cong \mathcal{B}_{\infty},$$

where the right-hand side is the inductive limit defined by the inductive system $\iota_{\bullet,\bullet}$ in (4.6).

Proof. The Toeplitz core $\mathcal{T}_{\mathfrak{H}}^{(0)}$ is the norm closure of linear combinations of elements of the form $S_{\mathbf{j}} S_{\mathbf{k}}^*$ with $|\mathbf{j}| = |\mathbf{k}|$ as well as their products with the vacuum projection $p_0 = |\Omega\rangle\langle\Omega|$. The elements which are products with $|\Omega\rangle\langle\Omega|$ belong to $\Gamma_0 = \mathcal{K} \cap \mathcal{T}_{\mathfrak{H}}^{(0)}$. Hence,

$$\mathcal{B}_{\infty} = \mathcal{T}_{\mathfrak{H}}^{(0)} / \Gamma_0 = \mathcal{T}_{\mathfrak{H}}^{(0)} / (\mathcal{K} \cap \mathcal{T}_{\mathfrak{H}}^{(0)}) = \mathcal{O}_{\mathfrak{H}}^{(0)},$$

as asserted. \square

Remark 4.13. The quasi-diagonal extension of \mathcal{B}_{∞} mentioned in Remark 4.6 can now be taken as $\mathcal{D} = \mathcal{T}_{\mathfrak{H}}^{(0)} + \mathcal{K}$ (cf. [Blac1, V.4.2.16]).

4.3 Cuntz–Pimsner algebras from inductive limits

For $m > 0$, let $\mathfrak{H}_{-m} := \overline{\mathfrak{H}_m}$ denote the conjugate Hilbert space of \mathfrak{H}_m . For all $k \in \mathbb{Z}$ we can consider the $\mathcal{B}(\mathfrak{H}_m)$ -module

$$\mathcal{E}_m^{(k)} := \mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+k}),$$

and the maps $\iota_{m,l}^{(k)} : \mathcal{E}_m^{(k)} \rightarrow \mathcal{E}_l^{(k)}$ defined by

$$\iota_{m,l}^{(k)}(X) := \sum_{|\mathbf{r}|=l-m} R_{\mathbf{r}} X R_{\mathbf{r}}^*|_{\mathfrak{H}_l}, \quad \forall X \in \mathcal{E}_m^{(k)}.$$

We define the C^* -algebras $\Gamma_b(\mathcal{E}_{\bullet}^{(k)})$ and $\Gamma_0(\mathcal{E}_{\bullet}^{(k)})$ in the same way as in Notation 4.1 and we denote by $\pi^{(k)} : \Gamma_b(\mathcal{E}_{\bullet}^{(k)}) \rightarrow \Gamma_b(\mathcal{E}_{\bullet}^{(k)})/\Gamma_0(\mathcal{E}_{\bullet}^{(k)})$ the quotient map.

Define ${}^0\mathcal{E}^{(k)}$ to be the vector space consisting of all elements of the form

$$\varsigma^{(m,k)}(X) := \pi^{(k)}(\iota_{m,l}^{(k)}(X))_{l \geq m}, \quad X \in \mathcal{E}_m^{(k)}$$

for all $m \in \mathbb{N}_0$. In particular, ${}^0\mathcal{E}^{(0)} \equiv {}^0\mathcal{B}_{\infty}$ is the algebraic part of $\mathcal{B}_{\infty} \equiv \mathcal{E}^{(0)} \cong \mathcal{O}_{\mathfrak{H}}^{(0)}$. Each ${}^0\mathcal{E}^{(k)}$ is a module over ${}^0\mathcal{B}_{\infty}$. The linear span of

$$\mathcal{B}_{\infty} {}^0\mathcal{E}^{(k)} := \{f\psi | f \in \mathcal{B}_{\infty}, \psi \in {}^0\mathcal{E}^{(k)}\}$$

is a module over \mathcal{B}_{∞} , which we denote by $\mathcal{E}^{(k)}$.

Theorem 4.14. *The Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ is isomorphic to the C^* -algebra generated by $\mathcal{E}^{(1)}$ and \mathcal{B}_{∞} . It allows the decomposition*

$$\mathcal{O}_{\mathfrak{H}} \cong \overline{\bigoplus_{k \in \mathbb{Z}} \mathcal{E}^{(k)}}^{\|\cdot\|}$$

and $\mathcal{E}^{(k)} \cong \mathcal{O}_{\mathfrak{H}}^{(k)}$ is the spectral subspace for the gauge action corresponding to $k \in \mathbb{Z}$.

Proof. The vector space $\mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+k})$ has an overcomplete basis given by the operators $S_{\mathbf{k}}|_{\mathfrak{H}_m}$ for all $\mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{k}| = k$. In particular, $\mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+1})$ is spanned by $S_j|_{\mathfrak{H}_m}$ for $j = 1, \dots, n$. Recalling that S_j is the shift by the basis vector $e_j \in \mathfrak{H}$, we see that

$$\begin{aligned} \iota_{m,l}^{(k)}(S_j|_{\mathfrak{H}_m}) &= \sum_{|\mathbf{r}|=l-m} R_{\mathbf{r}} S_j R_{\mathbf{r}}^*|_{\mathfrak{H}_l} \\ &= \sum_{|\mathbf{r}|=l-m} S_j R_{\mathbf{r}} R_{\mathbf{r}}^*|_{\mathfrak{H}_l} = S_j|_{\mathfrak{H}_l}. \end{aligned}$$

We can identify a sequence $X_{\bullet} = (X_m)_{m \in \mathbb{N}_0}$ of operators $X_m \in \mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+k})$ with an operator on Fock space $\mathfrak{H}_{\mathbb{N}}$. The effect of the quotient map $\pi^{(k)}$ on such a sequence X_{\bullet} is to take it to its image in the Calkin algebra $\mathcal{B}(\mathfrak{H}_{\mathbb{N}})/\mathcal{K}$. From (4.5) we therefore have (for $m \geq 1$)

$$\varsigma^{(m,1)}(S_j|_{\mathfrak{H}_m}) = \varsigma^{(1,1)}(S_j|_{\mathfrak{H}}) = \pi^{(1)}((S_j|_{\mathfrak{H}_m})_{m \in \mathbb{N}_0}) = \pi^{(1)}(S_j) = Z_j,$$

where Z_1, \dots, Z_n are the generators of $\mathcal{O}_{\mathfrak{H}}$. Similarly one gets that $\varsigma^{(l,k)}(S_{\mathbf{k}}|_{\mathfrak{H}_m})$ is just $Z_{\mathbf{k}}$ for all $\mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{k}| = k$ and all $l \geq m$. The adjoints $S_{\mathbf{k}}^*$ define elements of $\Gamma_b(\mathcal{E}_{\bullet}^{(-m)})$ for $|\mathbf{k}| = m$. So $\mathcal{E}^{(k)} \cong \mathcal{O}_{\mathfrak{H}}^{(k)}$ holds for all $k \in \mathbb{Z}$. \square

4.4 Formulas for covariant symbols

Our discussion about inductive limits associated to subproduct system has been based on shift operators on Fock space. We now observe that what we are doing is in fact a generalization of Berezin quantization. First we show that there is a very simple expression for the maps $\varsigma^{(m)}$.

Theorem 4.15. Let Z_1, \dots, Z_n be the images of the shifts of the subproduct system \mathfrak{H}_\bullet . Then the covariant symbol map $\varsigma^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{O}_{\mathfrak{H}}^{(0)}$ defined in (4.3) can be expressed as

$$\varsigma^{(m)}(A) = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} Z_{\mathbf{j}} Z_{\mathbf{k}}^*.$$

Proof. From (4.3) we see that we need to express ι_m in terms of the Toeplitz operators S_1, \dots, S_n ; applying π transforms these into Z_1, \dots, Z_n . But that is easily done using Lemma 4.9: the “second quantization” of A ,

$$\sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} S_{\mathbf{j}} S_{\mathbf{k}}^* \in \mathcal{B}(\mathfrak{H}_{\mathbb{N}}),$$

acts as $\iota_{m,l}(A)$ on \mathfrak{H}_l for $l \geq m$ and as 0 on \mathfrak{H}_l for $l < m$. Applying the quotient map π to it, we obtain

$$\pi\left(\sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} S_{\mathbf{j}} S_{\mathbf{k}}^*\right) = \pi((\iota_{m,l}(A))_{l \geq m}) = \varsigma^{(m)}(A),$$

and on the other hand,

$$\pi\left(\sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} S_{\mathbf{j}} S_{\mathbf{k}}^*\right) = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} Z_{\mathbf{j}} Z_{\mathbf{k}}^*.$$

□

Corollary 4.16. For all $\mathbf{j}, \mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{j}| = |\mathbf{k}| = m$ and all $l \geq m$ we have

$$\varsigma^{(l)}(S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_l}) = Z_{\mathbf{j}} Z_{\mathbf{k}}^*.$$

Example 4.17. Let \mathfrak{H}_\bullet be a commutative subproduct system and let M be the compact manifold it defines (see Corollary 3.7). Then $\varsigma^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{O}_{\mathfrak{H}}^{(0)}$ coincides with the Berezin covariant symbol map $\varsigma^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow C(M)$ (mentioned in §3.2).

Notation 4.18. Fix a faithful representation of $\mathcal{O}_{\mathfrak{H}}$ on a Hilbert space \mathcal{H} and let

$$u \in M_n(\mathbb{C}) \otimes \mathcal{B}(\mathcal{H})$$

be a unitary $n \times n$ matrix with values $u_{j,k} \in \mathcal{B}(\mathcal{H})$ such that the first row of u is given by the generators Z_1, \dots, Z_n of $\mathcal{O}_{\mathfrak{H}}$. Let u^c be the matrix obtained from u by taking adjoints of each entry $u_{j,k}$. Denote by u_m the restriction of $u^{\otimes m}$ from $\mathfrak{H}^{\otimes m}$ to \mathfrak{H}_m and similarly for u^c . Finally,

$$\alpha^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{B}(\mathfrak{H}_m) \otimes \mathcal{B}(\mathcal{H})$$

will be the map which takes $A \in \mathcal{B}(\mathfrak{H}_m)$ to $u_m(A \otimes \mathbf{1})u_m^*$.

The following formulas are known from the classical case to define the “Berezin covariant symbol” in case we quantize a coadjoint orbit $M = G/K$ (cf. [Per1], [Lan2]).

Proposition 4.19. Assume that $u^{\otimes m}$ preserves the subspace $\mathfrak{H}_m \subset \mathfrak{H}^{\otimes m}$, in the sense that $u_m = u^{\otimes m}(p_m \otimes \mathbf{1})$. Let $|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}|$ be the rank-1 projection onto the line spanned by $e_1^{\otimes m}$, where e_1 is the first basis vector in \mathfrak{H} . Then for all $A \in \mathcal{B}(\mathfrak{H}_m)$ we have

$$\varsigma^{(m)}(A) = (\text{Tr} \otimes \text{id})((A \otimes \mathbf{1})u_m^*(|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}| \otimes \mathbf{1})u_m^c) \quad (4.9)$$

$$= (\text{Tr} \otimes \text{id})(\alpha^{(m)}(A)(|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}| \otimes \mathbf{1})). \quad (4.10)$$

Proof. We have

$$u_m^c(|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}| \otimes \mathbf{1})u_m^c = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_m} \otimes Z_{\mathbf{k}} Z_{\mathbf{j}}^*$$

so (4.9) is clear. For (4.10) we can use the formula

$$\alpha^{(m)}(A) = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{r},\mathbf{s}} S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_m} \otimes u_{\mathbf{j},\mathbf{r}} u_{\mathbf{k},\mathbf{s}}^*.$$

□

4.5 Changing the inner products

Notation 4.20. From now on $Q \in \mathcal{B}(\mathfrak{H})$ is a positive invertible $n \times n$ matrix such that the operator $Q^{\otimes m}$ on $\mathfrak{H}^{\otimes m}$ preserves the subspace \mathfrak{H}_m for all $m \in \mathbb{N}_0$.

Saying that \mathfrak{H}_m is invariant under $Q^{\otimes m}$ is equivalent to saying that the compression

$$Q_m := p_m Q^{\otimes m} p_m$$

is equal to $Q^{\otimes m} p_m$.

We will use Q to define a new inner product. The role of Q may remain quite mysterious until we arrive at the quantum-group examples. In these examples, the subproduct system \mathfrak{H}_\bullet will itself depend on Q , in addition to the fact that Q defines the inner product. We can mention already that for the Berezin quantization of a *commutative* manifold (reviewed in §3) one should take $Q = p_1$, the identity operator on \mathfrak{H} .

Remark 4.21. The property $p_m Q^{\otimes m} p_m = Q^{\otimes m} p_m$ ensures that Q_m is invertible; its inverse is $p_m (Q^{\otimes m})^{-1} p_m$ because

$$(Q^{\otimes m})^{-1} p_m Q^{\otimes m} p_m = (Q^{\otimes m})^{-1} Q^{\otimes m} p_m = p_m,$$

$$Q^{\otimes m} p_m (Q^{\otimes m})^{-1} p_m = Q^{\otimes m} (Q^{\otimes m})^{-1} p_m = p_m,$$

where we used the fact that the inverse A^{-1} of any invertible matrix A preserves every A -invariant subspace. We denote by Q_m^{-1} this inverse of Q_m . For $l \geq m$ we have

$$Q_l^{-1} (Q_m \otimes Q_{l-m}) = (Q^{\otimes l})^{-1} ((Q^{\otimes m}) \otimes (Q^{\otimes (l-m)})) p_l = p_l,$$

where we regard p_l as an operator from $\mathfrak{H}_m \otimes \mathfrak{H}_{l-m}$ onto \mathfrak{H}_l .

We choose the orthonormal basis e_1, \dots, e_n such that Q is diagonal and, as before, we let S_1, \dots, S_n be the shifts on $\mathfrak{H}_\mathbb{N}$ by these basis vectors. We shall write

$$Q_{j,k} := (Q_m)_{j,k}, \quad Q^{j,k} := (Q_m^{-1})_{j,k}.$$

We associate to each Q_m a density matrix

$$\rho^{(m)} = \rho_Q^{(m)} := \frac{Q_m}{\text{Tr}(Q_m)},$$

and denote by ϕ_m the state on $\mathcal{B}(\mathfrak{H}_m)$ defined by

$$\phi_m(A) := \text{Tr}(\rho^{(m)} A), \quad \forall A \in \mathcal{B}(\mathfrak{H}_m).$$

From now on we shall assume that \mathfrak{H}_m is endowed with the $\rho^{(m)}$ -inner product.

$$\langle \psi | \phi \rangle_{\rho^{(m)}} := \langle \psi | \rho^{(m)} \phi \rangle, \quad \forall \phi, \psi \in \mathfrak{H}_m. \quad (4.11)$$

We stress that (unless $\mathfrak{H}_\bullet = \mathfrak{H}^{\otimes \bullet}$)

$$\rho^{(m)} \neq \frac{Q_m}{\text{Tr}(Q)^m} = p_m (\rho^{(1)})^{\otimes m} p_m.$$

4.6 The isometries

Now that \mathfrak{H}_m is endowed with the inner product (4.11) we discuss how $\mathcal{B}(\mathfrak{H}_m)$ can be mapped into $\mathcal{B}(\mathfrak{H}_l)$ when $m < l$ and calculate the explicit isometries $\mathfrak{H}_l \hookrightarrow \mathfrak{H}_{l-m} \otimes \mathfrak{H}_m$ and $\mathfrak{H}_l \hookrightarrow \mathfrak{H}_m \otimes \mathfrak{H}_{l-m}$. This construction would fail without the assumption that Q_m preserves \mathfrak{H}_m .

Proposition 4.22. *The isometry from \mathfrak{H}_l into $\mathfrak{H}_{l-m} \otimes \mathfrak{H}_m$ is given by*

$$\bar{V}_{m,l}\psi = \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}} \sum_{|\mathbf{r}|=l-m} p_l(e_{\mathbf{r}} \otimes S_{\mathbf{r}}^*\psi), \quad \forall \psi \in \mathfrak{H}_l, \quad (4.12)$$

and its adjoint by

$$\bar{V}_{m,l}^* = \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}} (\rho^{(l)})^{-1} (\rho^{(l-m)} \otimes \rho^{(m)}). \quad (4.13)$$

Proof. We proceed by first calculating the adjoint of the given operator (4.12). Let $\lambda_{m,l} := \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}}$. For all $\xi \in \mathfrak{H}_{l-m}, \eta \in \mathfrak{H}_m$ and all $\psi \in \mathfrak{H}_l$ we have

$$\begin{aligned} \langle \bar{V}_{m,l}^*(\xi \otimes \eta) | \psi \rangle_{\rho^{(l)}} &= \langle \xi \otimes \eta | \bar{V}_{m,l} \psi \rangle_{\rho^{(l-m)} \otimes \rho^{(m)}} \\ &= \lambda_{m,l} \sum_{|\mathbf{r}|=l-m} \langle \xi \otimes \eta | p_l(e_{\mathbf{r}} \otimes S_{\mathbf{r}}^*\psi) \rangle_{\rho^{(l-m)} \otimes \rho^{(m)}} \\ &= \lambda_{m,l} \sum_{|\mathbf{r}|=l-m} \langle \xi \otimes \eta | S_{\mathbf{r}} S_{\mathbf{r}}^* \psi \rangle_{\rho^{(l-m)} \otimes \rho^{(m)}} \\ &= \lambda_{m,l} \langle (\rho^{(l)})^{-1} (\rho^{(l-m)} \xi \otimes \rho^{(m)} \eta) | \psi \rangle_{\rho^{(l)}} \end{aligned}$$

and hence (4.13) holds. Finally, using Remark 4.21,

$$\begin{aligned} \bar{V}_{m,l}^* \bar{V}_{m,l} \psi &= \frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)} \sum_{|\mathbf{r}|=l-m} (\rho^{(l)})^{-1} (\rho^{(l-m)} e_{\mathbf{r}} \otimes \rho^{(m)} S_{\mathbf{r}}^* \psi) \\ &= \sum_{|\mathbf{r}|=l-m} Q_l^{-1} (Q_{l-m} e_{\mathbf{r}} \otimes Q_m S_{\mathbf{r}}^* \psi) = \sum_{|\mathbf{r}|=l-m} p_l(e_{\mathbf{r}} \otimes S_{\mathbf{r}}^* \psi) \\ &= \sum_{|\mathbf{r}|=l-m} S_{\mathbf{r}} S_{\mathbf{r}}^* \psi = \psi, \end{aligned}$$

so V is the desired isometry. Let us also calculate the final projection:

$$\bar{V}_{m,l} \bar{V}_{m,l}^* = \sum_{|\mathbf{r}|=l-m} S_{\mathbf{r}} S_{\mathbf{r}}^* Q_l^{-1} (Q_{l-m} \otimes Q_m) = p_l(p_{l-m} \otimes p_m).$$

□

Now let $A \in \mathcal{B}(\mathfrak{H}_l)$ and define

$$\bar{\iota}_{m,l}(A) := \bar{V}_{m,l}^*(\mathbf{1}_{\mathfrak{H}_{l-m}} \otimes A) \bar{V}_{m,l}. \quad (4.14)$$

We have

$$\begin{aligned} \bar{V}_{m,l}^*(\mathbf{1}_{\mathfrak{H}_{l-m}} \otimes A) \bar{V}_{m,l} \psi &= \frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)} \sum_{|\mathbf{r}|=l-m} (\rho^{(l)})^{-1} (\rho^{(l-m)} e_{\mathbf{r}} \otimes \rho^{(m)} A S_{\mathbf{r}}^* \psi) \\ &= \sum_{|\mathbf{r}|=l-m} Q_l^{-1} (Q_{l-m} e_{\mathbf{r}} \otimes Q_m A S_{\mathbf{r}}^* \psi) \\ &= \sum_{|\mathbf{r}|=l-m} p_l(e_{\mathbf{r}} \otimes A S_{\mathbf{r}}^* \psi) \\ &= \sum_{|\mathbf{r}|=l-m} S_{\mathbf{r}} A S_{\mathbf{r}}^* \psi, \end{aligned}$$

so $\bar{\iota}_{m,l}$ is a “chirality-flipped” version of $\iota_{m,l}$, i.e. the R_r ’s are replaced by S_j ’s (cf. (4.7)). If we use the inductive system $\bar{\iota}_{\bullet,\bullet}$ instead of $\iota_{\bullet,\bullet}$, the roles of the left and right Toeplitz algebras are interchanged.

Proposition 4.23. Define a coherent system of maps $\bar{\iota}_{m,l} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{B}(\mathfrak{H}_l)$ by

$$\bar{\iota}_{m,l}(A) := \bar{V}_{m,l}^*(\mathbf{1}_{\mathfrak{H}_{l-m}} \otimes A) \bar{V}_{m,l} \quad (4.15)$$

$$= \sum_{|\mathbf{s}|=l-m} S_{\mathbf{s}} A S_{\mathbf{s}}^*|_{\mathfrak{H}_l}. \quad (4.16)$$

Then $(\mathcal{B}(\mathfrak{H}_{\bullet}), \bar{\iota}_{\bullet,\bullet})$ is an inductive system, and the C^* -subalgebra of $\Gamma_b(\mathcal{B}_{\bullet})$ consisting of norm limits of the eventually constant sequences for $\bar{\iota}_{\bullet,\bullet}$ is equal to the $U(1)$ -invariant part of the right Toeplitz algebra $C^*(R_1, \dots, R_n)$.

Proof. The proof is very similar to the case of the $\iota_{m,l}$'s (the “left case”). \square

We furthermore note that if we expand $A \in \mathcal{B}(\mathfrak{H}_m)$ as $A = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} R_{\mathbf{j}} R_{\mathbf{k}}^*|_{\mathfrak{H}_m}$ then

$$\bar{\iota}_{m,l}(A) = \sum_{|\mathbf{j}|=m=|\mathbf{k}|} A_{\mathbf{j},\mathbf{k}} R_{\mathbf{j}} R_{\mathbf{k}}^*|_{\mathfrak{H}_l},$$

again similar to the left case. More will be said on the “chiral duality” between $\bar{\iota}_{\bullet,\bullet}$ and $\iota_{\bullet,\bullet}$ in Remark 5.7.

We now want to find an isometric implementation of $\iota_{m,l}$ similar to (4.14). For this we need to flip the tensor factors.

Proposition 4.24. The isometry from \mathfrak{H}_l into $\mathfrak{H}_m \otimes \mathfrak{H}_{l-m}$ is given by

$$V_{m,l}\psi = \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}} \sum_{|\mathbf{r}|=l-m} p_l(R_{\mathbf{r}}^* \psi \otimes e_{\mathbf{r}}), \quad \forall \psi \in \mathfrak{H}_l, \quad (4.17)$$

and its adjoint by

$$V_{m,l}^* = \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}} (\rho^{(l)})^{-1} (\rho^{(m)} \otimes \rho^{(l-m)}). \quad (4.18)$$

Proof. For all $\xi \in \mathfrak{H}_{l-m}$, $\eta \in \mathfrak{H}_m$ and all $\psi \in \mathfrak{H}_l$ we have

$$\begin{aligned} \langle V_{m,l}^*(\eta \otimes \xi) | \psi \rangle_{\rho^{(l)}} &= \langle \eta \otimes \xi | V_{m,l} \psi \rangle_{\rho^{(m)} \otimes \rho^{(l-m)}} \\ &= \lambda_{m,l} \sum_{|\mathbf{r}|=l-m} \langle \eta \otimes \xi | R_{\mathbf{r}} R_{\mathbf{r}}^* \psi \rangle_{\rho^{(m)} \otimes \rho^{(l-m)}} \\ &= \lambda_{m,l} \langle (\rho^{(l)})^{-1} (\rho^{(m)} \eta \otimes \rho^{(l-m)} \xi) | \psi \rangle_{\rho^{(l)}}, \end{aligned}$$

and the rest is similar to the proof of Proposition 4.24. \square

Corollary 4.25. The inductive system $\iota_{\bullet,\bullet}$ is implemented by the system $V_{\bullet,\bullet}$ of isometries:

$$\iota_{m,l}(A) = V_{m,l}^*(A \otimes \mathbf{1}_{\mathfrak{H}_{l-m}}) V_{m,l}, \quad \forall A \in \mathcal{B}(\mathfrak{H}_m).$$

Proof. Follows from formula (4.7) and the calculation (with $\psi \in \mathfrak{H}_l$)

$$\begin{aligned} V_{m,l}^*(A \otimes \mathbf{1}_{\mathfrak{H}_{l-m}}) V_{m,l} \psi &= \sqrt{\frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)}} \sum_{|\mathbf{r}|=l-m} (\rho^{(l)})^{-1} (\rho^{(m)} A R_{\mathbf{r}}^* \psi \otimes \rho^{(l-m)} e_{\mathbf{r}}) \\ &= \sum_{|\mathbf{r}|=l-m} Q_l^{-1} (Q_m A R_{\mathbf{r}}^* \psi \otimes Q_{l-m} e_{\mathbf{r}}) \\ &= \sum_{|\mathbf{r}|=l-m} p_l(A R_{\mathbf{r}}^* \psi \otimes e_{\mathbf{r}}) \\ &= \sum_{|\mathbf{r}|=l-m} R_{\mathbf{r}} A R_{\mathbf{r}}^* \psi. \end{aligned}$$

\square

5 Projective limits

We now want to realize the same algebra $\mathcal{O}_{\mathfrak{H}}^{(0)}$ as a projective limit. For this we need some background information from [Hawk1, §B2].

5.1 Relaxing the notion of projective limit

In this paper, a “projective limit” will always refer to the following object which, in comparison to more conventional C^* -algebraic projective limits, is defined in terms of completely positive maps instead of C^* -homomorphisms.

Definition 5.1 ([Hawk1, §B2]). A **projective system** $(\mathcal{B}_\bullet, j_{\bullet, \bullet})$ is a sequence of finite-dimensional matrix algebras \mathcal{B}_m and norm-contracting completely positive mappings $j_{l, m} : \mathcal{B}_l \rightarrow \mathcal{B}_m$ for $m \leq l$ satisfying $j_{l, m} = j_{r, m} \circ j_{l, r}$ for all $m \leq r \leq l$. The **projective limit** of $(\mathcal{B}_\bullet, j_{\bullet, \bullet})$ is the vector space defined by

$$\mathcal{B}^\infty := \{A_\bullet = (A_m)_{m \in \mathbb{N}_0} \in \Gamma_b(\mathcal{B}_\bullet) \mid A_{m-1} = j_{m, m-1}(A_m) \text{ for all } m \in \mathbb{N}\},$$

equipped with the norm

$$\|A_\bullet\| := \lim_{m \rightarrow \infty} \|A_m\|.$$

Remark 5.2. The intersection of \mathcal{B}^∞ with Γ_0 is $\{0\}$. We always identify \mathcal{B}^∞ with its embedding into $\Gamma_b(\mathcal{B}_\bullet)/\Gamma_0(\mathcal{B}_\bullet)$ because it is more likely that \mathcal{B}^∞ is an algebra when multiplication is taken modulo Γ_0 . If we do so and then pull back \mathcal{B}^∞ via the quotient map $\pi : \Gamma_b \rightarrow \Gamma_b/\Gamma_0$, we obtain a vector space $\pi^{-1}(\mathcal{B}^\infty)$ which is much larger than \mathcal{B}^∞ , namely

$$\pi^{-1}(\mathcal{B}^\infty) = \mathcal{B}^\infty \cup \Gamma_0(\mathcal{B}_\bullet). \quad (5.1)$$

Importantly, \mathcal{B}^∞ is an algebra (hence a C^* -algebra) if and only if (5.1) is.

Remark 5.3. We could also define \mathcal{B}^∞ as the set of elements

$$f = \pi((j_{\infty, m}(f))_{m \in \mathbb{N}_0})$$

where the components $j_{\infty, m}(f) \in \mathcal{B}_m$ satisfy $j_{\infty, m}(f) = j_{l, m} \circ j_{\infty, l}(f)$. We can regard $j_{\infty, m} := \lim_{l \rightarrow \infty} j_{l, m}$ as the map from \mathcal{B}^∞ to \mathcal{B}_m which evaluates $A_\bullet = (A_m)_{m \in \mathbb{N}_0} \in \mathcal{B}^\infty$ at m ,

$$j_{\infty, m}(A_\bullet) = A_m.$$

5.2 Projective system for subproduct systems

Let \mathfrak{H}_\bullet be a subproduct system. We let $Q \in \mathcal{B}(\mathfrak{H})$ be as in Notation 4.20 and endow \mathfrak{H}_m with the inner product defined by the density matrix $\rho^{(m)} := Q_m / \text{Tr}(Q_m)$.

Lemma 5.4. Define maps $j_{l, m} : \mathcal{B}(\mathfrak{H}_l) \rightarrow \mathcal{B}(\mathfrak{H}_m)$ for $m \leq l$ by

$$j_{l, m}(A) := \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|\mathbf{k}|=l-m} (Q^{\otimes m})_{\mathbf{k}, \mathbf{k}} R_{\mathbf{k}}^* A R_{\mathbf{k}}|_{\mathfrak{H}_m}, \quad \forall A \in \mathcal{B}(\mathfrak{H}_l). \quad (5.2)$$

Then, with $V_{m, l}$ as in Proposition 4.24, we have the formula

$$j_{l, m}(A) = (\text{id}_{\mathcal{B}_m} \otimes \phi_{l-m})(V_{m, l} A V_{m, l}^*) \quad (5.3)$$

and $j_{\bullet, \bullet}$ is a projective system.

Proof. First of all, for all $A \in \mathcal{B}(\mathfrak{H}_l)$ we have

$$\begin{aligned} j_{r,m} \circ j_{l,r}(A) &= \frac{\text{Tr}(Q_r)}{\text{Tr}(Q_l)} \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_r)} \sum_{|j|=r-m, |k|=l-r} (Q^{\otimes m})_{j,j} (Q^{\otimes m})_{k,k} R_j^* R_k^* A R_k R_j|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|kj|=l-m} (Q^{\otimes m})_{kj,kj} R_{kj}^* A R_{kj}|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|r|=l-m} (Q^{\otimes m})_{r,r} R_r^* A R_r|_{\mathfrak{H}_m} = j_{l,m}(A), \end{aligned}$$

and it is obvious that each $j_{l,m}$ is completely positive. The norm-contracting property holds because $j_{l,m}$ is in fact unital. To see this we first prove the alternative formula (5.3). We have

$$\begin{aligned} j_{l,m}(A) &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|r|=l-m} (Q^{\otimes m})_{r,r} R_r^* A R_r|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|r|=l-m} \sum_{|j|=m=|k|} (Q^{\otimes m})_{r,r} \langle e_j | R_r^* A R_r e_k \rangle S_j S_k^*|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|r|=l-m} \sum_{|j|=m=|k|} (Q^{\otimes m})_{r,r} \langle e_j \otimes e_r | A (e_k \otimes e_r) \rangle S_j S_k^*|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|r|=l-m} \sum_{|j|=m=|k|} (Q^{\otimes m})_{r,r} (A)_{jr,kr} S_j S_k^*|_{\mathfrak{H}_m} \\ &= (\text{id} \otimes \phi_{l-m})(V_{m,l} A V_{m,l}^*), \end{aligned}$$

where in the last equality we used that, for all $\xi_1, \xi_2 \in \mathfrak{H}_m$, $\eta_1, \eta_2 \in \mathfrak{H}_{l-m}$,

$$\begin{aligned} &\langle V_{m,l}^*(\xi_1 \otimes \eta_1) | A V_{m,l}^*(\xi_2 \otimes \eta_2) \rangle \\ &= \frac{\text{Tr}(Q_m) \text{Tr}(Q_{l-m})}{\text{Tr}(Q_l)} \langle \rho^{(l)-1}(\rho^{(m)} \xi_1 \otimes \rho^{(l-m)} \eta_1) | A \rho^{(l)-1}(\rho^{(m)} \xi_2 \otimes \rho^{(l-m)} \eta_2) \rangle \\ &= \langle \xi_1 \otimes \eta_1 | A \rho^{(l)-1}(\rho^{(m)} \xi_2 \otimes \rho^{(l-m)} \eta_2) \rangle \\ &= \frac{\text{Tr}(Q_l)}{\text{Tr}(Q_{l-m}) \text{Tr}(Q_m)} \langle \xi_1 \otimes \eta_1 | A (\xi_2 \otimes \eta_2) \rangle, \end{aligned}$$

so that summing such inner products over a basis for $\mathfrak{H}_m \otimes \mathfrak{H}_{l-m}$ and multiplying with $\frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} S_j S_k^*|_{\mathfrak{H}_m}$ is the same thing as partially tracing $V_{m,l} A V_{m,l}^*$ with $Q_{l-m} / \text{Tr}(Q_{l-m})$.

Now (5.3) gives

$$j_{l,m}(p_l) = (\text{id}_{\mathcal{B}_m} \otimes \phi_{l-m})(p_m \otimes p_{l-m}) p_l (p_m \otimes p_{l-m}) = \phi_{l-m}(p_{l-m}) p_m = p_m,$$

showing that $j_{l,m}$ is unital. □

5.3 The state on $\mathcal{O}_{\mathfrak{H}}$

5.3.1 $j_{l,m}$ as adjoint of $\iota_{m,l}$

Proposition 5.5. *Let $m, l \in \mathbb{N}_0$ with $m \leq l$. Then $j_{l,m}$ is the adjoint of $\iota_{m,l}$: for all $A \in \mathcal{B}(\mathfrak{H}_l)$ and all $B \in \mathcal{B}(\mathfrak{H}_m)$ we have*

$$\phi_l(A \iota_{m,l}(B)) = \phi_m(j_{l,m}(A) B). \quad (5.4)$$

In particular, taking $A = p_l$ respectively $B = p_m$ we obtain the compatibility relations

$$\phi_l \circ \iota_{m,l} = \phi_m, \quad (5.5)$$

$$\phi_l = \phi_m \circ j_{l,m}. \quad (5.6)$$

Proof. We have

$$\begin{aligned}
\phi_l(A\iota_{m,l}(B)) &= \phi_l(AV_{m,l}^*(B \otimes p_{l-m})V_{m,l}) \\
&= (\phi_m \otimes \phi_{l-m})(V_{m,l}AV_{m,l}^*(B \otimes p_{l-m})) \\
&= \phi_m((\text{id}_{\mathcal{B}_m} \otimes \phi_{l-m})(V_{m,l}AV_{m,l}^*(B \otimes p_{l-m}))) \\
&= \phi_m((\text{id}_{\mathcal{B}_m} \otimes \phi_{l-m})(V_{m,l}AV_{m,l}^*B)),
\end{aligned}$$

which equals $\phi_m(j_{l,m}(A)B)$ by (5.3). \square

Corollary 5.6. *The states ϕ_m satisfy the “right invariance” condition*

$$\phi_m(j_{l,m} \circ \iota_{m,l}(A)) = \phi_m(A), \quad \forall A \in \mathcal{B}(\mathfrak{H}_m). \quad (5.7)$$

Proof. Just use (5.6) and then (5.5). \square

Remark 5.7. Similarly one shows that

$$\phi_m = \phi_l \circ \bar{\iota}_{m,l}. \quad (5.8)$$

for the “right” inductive system $\bar{\iota}_{\bullet,\bullet}$. The adjoint of $\bar{\iota}_{m,l}$ is

$$\bar{j}_{l,m}(A) = \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|\mathbf{r}|=l-m} (Q^{\otimes m})_{\mathbf{r},\mathbf{r}} S_{\mathbf{r}}^* A S_{\mathbf{r}}|_{\mathfrak{H}_m} = (\phi_{l-m} \otimes \text{id})(\bar{V}_{m,l} A \bar{V}_{m,l}^*).$$

A “left invariance” condition similar to (5.7) is also deduced using the $\bar{\iota}_{m,l}$ ’s and their adjoints.

5.3.2 The limit state

Corollary 5.8. *The limit*

$$\phi_\infty := \lim_{m \rightarrow \infty} \phi_m$$

is a well-defined state on $\mathcal{T}_{\mathfrak{H}}^{(0)}$. It annihilates $\mathcal{T}_{\mathfrak{H}}^{(0)} \cap \mathcal{K}$, so it descends to a state ω_Q on $\mathcal{O}_{\mathfrak{H}}^{(0)} = \mathcal{B}_\infty$.

Proof. For well-definedness we use (5.5) and recall that the elements of $\mathcal{T}_{\mathfrak{H}}^{(0)}$ are norm limits of eventually constant under $\iota_{\bullet,\bullet}$. The fact that ϕ_∞ descends to $\mathcal{T}_{\mathfrak{H}}^{(0)}/(\mathcal{T}_{\mathfrak{H}}^{(0)} \cap \mathcal{K})$ follows from

$$|\phi_m(A)| \leq \|A\|, \quad \forall A \in \mathcal{B}(\mathfrak{H}_m), m \in \mathbb{N}_0,$$

since this shows that $\lim_m \|A_m\| = 0$ implies $\phi_\infty(A_\bullet) = 0$ for all $A_\bullet = (A_m)_{m \in \mathbb{N}_0} \in \mathcal{T}_{\mathfrak{H}}^{(0)}$. \square

Proposition 5.9. *The state $\omega_Q : \mathcal{B}_\infty \rightarrow \mathbb{C}$ is KMS, with modular automorphism group $\sigma_\bullet = (\sigma_t)_{t \in \mathbb{R}}$ given by*

$$\sigma_t \circ \varsigma^{(m)}(A) = \varsigma^{(m)}(Q_m^{it} A Q_m^{-it})$$

for all $A \in \mathcal{B}(\mathfrak{H}_m)$ and all $m \in \mathbb{N}_0$, and ω_Q satisfies

$$\omega_Q(Z_{\mathbf{j}} Z_{\mathbf{k}}^*) = \frac{Q_{\mathbf{k},\mathbf{j}}}{\text{Tr}(Q_m)} \quad (5.9)$$

for all $\mathbf{j}, \mathbf{k} \in \mathbb{F}_n^+$ with $|\mathbf{j}| = |\mathbf{k}| = m$. Moreover, the covariant symbol map $\varsigma^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{B}_\infty$ intertwines ω_Q and ϕ_m :

$$\omega_Q \circ \varsigma^{(m)} = \phi_m. \quad (5.10)$$

Proof. Due to (5.5) we have, if $|\mathbf{j}| = |\mathbf{k}| = m$,

$$\begin{aligned}\omega_Q(Z_{\mathbf{j}}Z_{\mathbf{k}}^*) &= \phi_\infty(S_{\mathbf{j}}S_{\mathbf{k}}^*) \\ &= \lim_{m \leq l \rightarrow \infty} \phi_l(S_{\mathbf{j}}S_{\mathbf{k}}^*p_l) \\ &= \phi_m(S_{\mathbf{j}}S_{\mathbf{k}}^*p_m),\end{aligned}$$

and so the first formula in (5.9) follows from

$$\begin{aligned}\mathrm{Tr}(Q_m)\phi_m(S_{\mathbf{j}}S_{\mathbf{k}}^*) &= \sum_{|\mathbf{r}|=m} (Q^{\otimes m})_{\mathbf{r},\mathbf{r}} \langle e_{\mathbf{r}} | p_m e_{\mathbf{j}} \rangle \langle e_{\mathbf{k}} | p_m e_{\mathbf{r}} \rangle \\ &= \langle e_{\mathbf{k}} | Q_m e_{\mathbf{j}} \rangle.\end{aligned}$$

The definition of $\zeta^{(m)}$ immediately gives (5.10), again using (5.5).

That ω_Q is KMS follows from (5.10), in view of the fact that the $*$ -algebra generated by the covariant symbols $\zeta^{(m)}(A)$ is dense in \mathcal{B}_∞ and that each ϕ_m is KMS. Finally, for $t \in \mathbb{R}$ the modular automorphism $\sigma_t^{\phi_m}$ of ϕ_m takes $A \in \mathcal{B}(\mathfrak{H}_m)$ to $(\rho^{(m)})^{it} A (\rho^{(m)})^{-it} = Q_m^{it} A Q_m^{-it}$. \square

We can extend ω_Q to a state, still denoted by ω_Q , on the whole Cuntz–Pimsner algebra by defining it to be zero on each spectral subspace $\mathcal{O}_{\mathfrak{H}}^{(k)}$ except $\mathcal{O}_{\mathfrak{H}}^{(0)}$.

Remark 5.10. From now on we shall assume that ω_Q is faithful. In general, we could go the GNS representation of \mathcal{B}_∞ associated with ω_Q and use the faithful state induced by ω_Q on the image of \mathcal{B}_∞ , which is a quotient of \mathcal{B}_∞ (recall that ω_Q is KMS). Then our results hold for the image of \mathcal{B}_∞ in the GNS representation.

Example 5.11. For the product system $\mathfrak{H}^{\otimes \bullet}$, the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ is the Cuntz algebra \mathcal{O}_n and ω_Q is the quasi-free state on \mathcal{O}_n defined by the density matrix $Q/\mathrm{Tr}(Q)$ [Ev1].

5.4 Contravariant symbols

Proposition 5.12. *The adjoint $\zeta^{(m)} : \mathcal{B}_\infty \rightarrow \mathcal{B}(\mathfrak{H}_m)$ of the covariant symbol map $\zeta^{(m)} : \mathcal{B}(\mathfrak{H}_m) \rightarrow \mathcal{B}_\infty$, defined by the relation*

$$\omega_Q(\zeta^{(m)}(A)^* f) = \phi_m(A^* \zeta^{(m)}(f)), \quad \forall A \in \mathcal{B}(\mathfrak{H}_m), f \in \mathcal{B}_\infty,$$

is given by

$$\zeta^{(m)}(f) = \mathrm{Tr}(Q_m) \sum_{|\mathbf{j}|, |\mathbf{k}|=m} (Q^{\otimes m})_{\mathbf{j},\mathbf{j}}^{-1} \omega_Q(Z_{\mathbf{j}}Z_{\mathbf{k}}^* f) S_{\mathbf{k}} S_{\mathbf{j}}^*|_{\mathfrak{H}_m} \quad (5.11)$$

Proof. Let $\zeta^{(m)}(f)$ be defined by (5.11). Then

$$\begin{aligned}\omega_Q(\zeta^{(m)}(A)^* f) &= \sum_{|\mathbf{j}|=m=|\mathbf{k}|} (A^*)_{\mathbf{j},\mathbf{k}} \omega_Q(Z_{\mathbf{j}}Z_{\mathbf{k}}^* f) \\ &= \sum_{|\mathbf{j}|=m=|\mathbf{k}|} (Q^{\otimes m})_{\mathbf{j},\mathbf{j}} (Q^{\otimes m})_{\mathbf{j},\mathbf{j}}^{-1} (A^*)_{\mathbf{j},\mathbf{k}} \omega_Q(Z_{\mathbf{j}}Z_{\mathbf{k}}^* f) \\ &= \phi_m(A^* \zeta^{(m)}(f)).\end{aligned}$$

\square

Corollary 5.13. *We have*

$$\omega_Q = \phi_m \circ \zeta^{(m)} \quad (5.12)$$

and, moreover, each $\zeta^{(m)}$ is unital.

Proof. Equation (5.12) is a direct consequence of the fact that $\zeta^{(m)}$ is adjoint to the unital map $\varsigma^{(m)}$. Unitality of $\zeta^{(m)}$ follows from $\omega_Q(\varsigma^{(m)}(A)^*\mathbf{1}) = \phi_m(A^*)$, which we know from (5.10). \square

We can now assembly the $\zeta^{(m)}$'s to a map

$$\zeta := \prod_{m \in \mathbb{N}_0} \zeta^{(m)} : \mathcal{B}_\infty \rightarrow \prod_{m \in \mathbb{N}_0} \mathcal{B}(\mathfrak{H}_m), \quad (5.13)$$

which is a noncommutative generalization of the total Toeplitz map (3.5). We can recover its components as

$$\zeta^{(m)}(f) = \zeta(f)p_m.$$

It is the following result which relies on the assumption that ω_Q is faithful (cf. Remark 5.10).

Lemma 5.14. *No nonzero element of \mathcal{B}_∞ is mapped to $\Gamma_0 = \mathcal{T}_{\mathfrak{H}}^{(0)} \cap \mathcal{K}$ under the map ζ .*

Proof. We have $\phi_m(\zeta^{(m)}(f)) = \omega_Q(f)$, so if $\zeta^{(m)}(f) \rightarrow 0$ as $m \rightarrow \infty$ then $\omega_Q(f) = 0$. Hence if $f \geq 0$ then $f = 0$ and the result follows. \square

Let $\mathcal{M} = \pi_{\omega_Q}(\mathcal{B})''$ be the von Neumann algebra generated by the inductive limit \mathcal{B}_∞ in the GNS representation of the limit state ω_Q . Then we can define $\zeta(f) \in \Gamma_b$ also for elements in \mathcal{M} , and Lemma 5.14 extends to \mathcal{M} .

Lemma 5.15. *For all $f \in \mathcal{M}$ and all $l \geq m$,*

$$\zeta^{(m)}(f) = j_{l,m} \circ \zeta^{(l)}(f). \quad (5.14)$$

Hence the image of \mathcal{M} under the total Toeplitz map ζ is contained in the projective limit \mathcal{B}^∞ , and in fact we have equality

$$\zeta(\mathcal{M}) = \mathcal{B}^\infty.$$

Therefore \mathcal{B}^∞ can be identified with the weak-closed operator system of elements of the form*

$$\zeta(f) = (\zeta^{(m)}(f))_{m \in \mathbb{N}_0}, \quad f \in \mathcal{M}$$

and, as in Remark 5.3,

$$\zeta^{(m)} = j_{\infty,m}$$

is the map which evaluates $(X_m)_{m \in \mathbb{N}_0} \in \mathcal{B}^\infty$ at $m \in \mathbb{N}_0$. The norm-closed subset $\zeta(\mathcal{B}_\infty)$ equals the anti-normally ordered part of the Toeplitz core.

Proof. We know that ζ is injective (Lemma 5.14). Since we have shown that $j_{l,m}$ is adjoint to $\iota_{m,l}$, we obtain (5.14) by taking adjoints of

$$\zeta^{(l)} \circ \iota_{m,l} = \zeta^{(m)}.$$

From (5.14) follows that $\zeta(f) \in \mathcal{B}^\infty$ for all $f \in \mathcal{M}$. Moreover, $\zeta : \mathcal{M} \rightarrow \Gamma_b$ is onto \mathcal{B}^∞ because each $\zeta^{(m)}$ is onto. Thus \mathcal{B}^∞ is in bijection with \mathcal{M} via ζ .

We need to show that $\zeta(\mathcal{B}_\infty)$ equals the anti-normally part of the Toeplitz core $\mathcal{T}_{\mathfrak{H}}^{(0)}$. Firstly, since the left and right shifts commute outside the vacuum subspace, for all $r, s = 1, \dots, n$ we have

$$\begin{aligned} j_{l,m}(S_r^* S_s p_l) &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|\mathbf{k}|=l-m} (Q^{\otimes m})_{\mathbf{k}, \mathbf{k}} R_{\mathbf{k}}^* S_r^* S_s R_{\mathbf{k}}|_{\mathfrak{H}_m} \\ &= \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|\mathbf{k}|=l-m} (Q^{\otimes m})_{\mathbf{k}, \mathbf{k}} S_r^* R_{\mathbf{k}}^* R_{\mathbf{k}} S_s|_{\mathfrak{H}_m} = S_r^* S_s|_{\mathfrak{H}_m} \end{aligned}$$

(where we used that $j_{l,m}(p_l) = p_m$), which shows that the anti-normally ordered elements of $\mathcal{T}_{\mathfrak{H}}^{(0)}$ are constant under $j_{\bullet, \bullet}$. Secondly, an explicit calculation using (5.11) shows that $\zeta(f)$ is anti-normally ordered for each $f \in \mathcal{B}_\infty$. \square

Remark 5.16. Now we can give an alternative proof for the fact that the contravariant symbol map $\zeta^{(l)} : \mathcal{B}_\infty \rightarrow \mathcal{B}(\mathfrak{H}_m)$ intertwines ω_Q with ϕ_l ,

$$\omega_Q = \phi_l \circ \zeta^{(l)}.$$

Recall that ω_Q denotes the limit state $\phi_\infty := \lim_{m \rightarrow \infty} \phi_m$ when regarded as a state on the quotient \mathcal{B}_∞ of $\mathcal{T}_{\mathfrak{H}}^{(0)}$. Then $\omega_Q = \phi_l \circ \zeta^{(l)}$ follows from the compatibility $\phi_m = \phi_l \circ j_{m,l}$ (see (5.5)) and the fact that $j_{\infty,l} = \zeta^{(l)}$.

5.5 The asymptotic multiplication

We now endow the projective limit \mathcal{B}^∞ with a multiplication which is the $m \rightarrow \infty$ limit of the multiplication on $\mathcal{B}(\mathfrak{H}_m)$.

Definition 5.17. The **projective-limit multiplication** on $\mathcal{B}^\infty \subset \Gamma_b$ is defined by

$$\zeta(f) \cdot \zeta(g) := \lim_{m \rightarrow \infty} \zeta^{(m)}(fg) \quad (5.15)$$

for all $f, g \in \mathcal{B}_\infty$.

The projective limit \mathcal{B}^∞ is not an algebra under the projective-limit multiplication, but we shall see that the subset $\zeta(\mathcal{B}_\infty)$ is.

The multiplication on \mathcal{B}^∞ taken modulo Γ_0 is the one where sequences $(\zeta^{(m)}(f))_{m \in \mathbb{N}_0}$ and $(\zeta^{(m)}(g))_{m \in \mathbb{N}_0}$ are multiplied componentwise but the finite- m part is ignored. That is,

$$\pi(\zeta(f)\zeta(g)) = \lim_{m \rightarrow \infty} \zeta^{(m)}(f)\zeta^{(m)}(g). \quad (5.16)$$

We will see momentarily that the products (5.15) and (5.16) coincide for $f, g \in \mathcal{B}_\infty$. Comparing the two formulas one then concludes that the Toeplitz maps $\zeta^{(m)}$ are “asymptotically multiplicative”. Again the projective limit \mathcal{B}^∞ is not an algebra under the multiplication modulo compacts, while $\zeta(\mathcal{B}_\infty)$ will be shown to be so.

Remark 5.18 (Filters). A projective-limit multiplication can be defined using any filter ω on \mathbb{N} . On the C^* -level this corresponds to considering not a subalgebra of Γ_b/Γ_0 but a subalgebra of Γ_b/Γ_ω where Γ_ω is the ideal consisting of the sequences A_\bullet with

$$\lim_{\omega} \|A_m\| = 0.$$

We recover Γ_0 if ω is the free filter of all cofinite subsets of \mathbb{N}_0 . Confer [RoSt1, §6.2].

5.6 The adjoint of the total Toeplitz map

Lemma 5.19. For $A \in \mathcal{B}(\mathfrak{H}_l)$ we have

$$j_{l,0}(A) = \phi_l(A)p_0. \quad (5.17)$$

and hence, if $\hat{\varepsilon}$ denotes the vacuum state restricted to $\mathcal{T}_{\mathfrak{H}}^{(0)}$,

$$\hat{\varepsilon} \circ j_{l,0} = \phi_l.$$

The vacuum state $\hat{\varepsilon}$ restricted to \mathcal{B}^∞ is equal to $\hat{\varepsilon} \circ j_{\infty,0}$ and coincides with the limit state $\phi_\infty := \lim_{l \rightarrow \infty} \phi_l$,

$$\hat{\varepsilon} \circ j_{\infty,0} = \phi_\infty.$$

Proof. We use $\phi_m \circ j_{l,m} = \phi_l$ for $m = 0$. This gives (5.17). The rest is obvious. \square

We can therefore regard $\hat{\varepsilon}$ as a state on the projective limit modulo compact operators as well, i.e. on the algebra $\pi(\mathcal{B}^\infty) \subset \Gamma_b/\Gamma_0$.

Recall that the covariant-symbol map $\zeta^{(m)}$ is the adjoint of $\zeta^{(m)}$, for each $m \in \mathbb{N}$. We now show that the total Toeplitz map ζ has an adjoint as well. This should be compared with [INT1, Lemma 2.3].

Proposition 5.20. *There exists a completely positive map*

$$\zeta^* : \zeta(\mathcal{B}_\infty) \rightarrow \mathcal{B}_\infty$$

such that, for all $X \in \zeta(\mathcal{B}_\infty)$ and all $f \in \mathcal{B}_\infty$,

$$\omega_Q(\zeta^*(X^*)f) = \hat{\varepsilon}(\pi(X^*)\pi(\zeta(f))). \quad (5.18)$$

Explicitly, this map is given by the point-norm limit $\zeta^* = \lim_{m \rightarrow \infty} \zeta^{(m)}$,

$$\zeta^*(X) = \lim_{m \rightarrow \infty} \zeta^{(m)}(Xp_m), \quad \forall X \in \zeta(\mathcal{B}_\infty),$$

and will be denoted by ς .

Proof. We identify $X \in \zeta(\mathcal{B}_\infty)$ with a bounded sequence $(X_m)_{m \in \mathbb{N}_0}$ of operators $X_m = Xp_m \in \mathcal{B}(\mathfrak{H}_m)$. Using the formula (5.16) for the multiplication in $\pi(\mathcal{B}^\infty)$ we have, by norm-continuity of the vacuum state, the norm limits

$$\begin{aligned} \hat{\varepsilon}(\pi(X^*)\pi(\zeta(f)))p_0 &= \langle \Omega | \lim_{m \rightarrow \infty} X_m^* \zeta^{(m)}(f) \Omega \rangle p_0 \\ &= J_{\infty,0} \left(\lim_{m \rightarrow \infty} X_m^* \zeta^{(m)}(f) \right) \\ &= \lim_{m \rightarrow \infty} J_{m,0} (X_m^* \zeta^{(m)}(f)) \\ &= \lim_{m \rightarrow \infty} \phi_m (X_m^* \zeta^{(m)}(f)) p_0 \\ &= \lim_{m \rightarrow \infty} \omega_Q (\zeta^{(m)}(X_m^*)f) p_0 \\ &= \omega_Q \left(\lim_{m \rightarrow \infty} \zeta^{(m)}(X_m^*)f \right) p_0. \end{aligned}$$

Being a point-norm limit of completely positive maps, ς is completely positive. \square

Since ζ is an isometry, ς is also the inverse of ζ , and we have

$$\text{id} = \varsigma \circ \zeta = \lim_{m \rightarrow \infty} \zeta^{(m)} \circ \zeta^{(m)},$$

making Remark 4.7 explicit. We have now seen that $\zeta : \mathcal{B}_\infty \rightarrow \zeta(\mathcal{B}_\infty)$ is a complete order isomorphism, i.e. a bijective unital completely positive map with completely positive inverse.

There is also a version of this result on the level of von Neumann algebras. As we shall see in 6.5.2, for any subproduct system \mathfrak{H}_\bullet , the weak- $*$ -closed operator system \mathcal{B}^∞ becomes a von Neumann algebra when equipped with a SOT-version of the projective-limit multiplication (5.15). When \mathfrak{H}_\bullet is the \mathbb{G} -subproduct system (see §6 below), \mathcal{B}^∞ is an operator system in the group-von Neumann algebra $\mathcal{R}(\mathbb{G})$.

Corollary 5.21. *The total Toeplitz map ζ intertwines the state ω_Q on $\mathcal{M} = \pi_{\omega_Q}(\mathcal{B}_\infty)''$ with the vacuum state $\hat{\varepsilon}$ on $\mathcal{B}^\infty \subset \mathcal{B}(\mathfrak{H}_\mathbb{N})$,*

$$\omega_Q = \hat{\varepsilon} \circ \zeta, \quad (5.19)$$

and similarly

$$\omega_Q \circ \varsigma = \hat{\varepsilon}, \quad (5.20)$$

Proof. Take $X = \mathbf{1}$ respectively $f = \mathbf{1}$ in (5.18) to get (5.19) respectively (5.20). \square

5.7 $\mathcal{O}_{\mathfrak{H}}^{(0)}$ as a projective limit

Theorem 5.22. *The operator system $\zeta(\mathcal{B}_\infty) \subset \mathcal{B}^\infty$ is a C^* -algebra with the multiplication taken modulo compacts; indeed $\pi(\zeta(\mathcal{B}_\infty))$ is isomorphic as a C^* -algebra to the inductive limit $\mathcal{B}_\infty \cong \mathcal{O}_{\mathfrak{H}}^{(0)}$.*

Remark 5.23. We saw in Lemma 5.14 that the image of ζ does not contain Γ_0 . On the other hand, $\zeta(\mathcal{B}_\infty) + \Gamma_0$ is generated by $\text{Ran } \zeta$ alone as a C^* -algebra. The theorem implies that

$$\zeta(f)\zeta(g) - \zeta(fg) \in \Gamma_0$$

even for nontrivial $f, g \in \mathcal{B}_\infty$. These elements do not belong to \mathcal{B}^∞ , which is why we need to apply the quotient map $\pi : \Gamma_b \rightarrow \Gamma_b/\Gamma_0$ in order to obtain an algebra.

Proof. This is a well-known consequence of the fact that $\zeta : \mathcal{B}_\infty \rightarrow \zeta(\mathcal{B}_\infty)$ is a complete order isomorphism from a C^* -algebra \mathcal{B}_∞ onto the operator system $\zeta(\mathcal{B}_\infty)$; see [Ar10, Prop. 2.2]. \square

Corollary 5.24. *The projective-limit multiplication on $\zeta(\mathcal{B}_\infty) \subset \Gamma_b$ coincides with the multiplication on $\zeta(\mathcal{B}_\infty)$ taken modulo Γ_0 .*

Proof. By uniqueness of the C^* -algebraic structure we know that any two multiplications on $\zeta(\mathcal{B}_\infty)$ compatible with the norm must be isomorphic. But as remarked in §5.5, the *exact* equality of the two products at hand is equivalent to the statement that ζ is multiplicative modulo Γ_0 , whence the result. \square

Now let us return to the problem of normal ordering in the Toeplitz algebra.

Proof of Lemma 2.15 (iii). The statement we want to prove is $\mathcal{T}_{\mathfrak{H}} = \overline{\text{span}}(\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}} \cup \mathcal{K})$, where $\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}}$ is the anti-normally ordered part of the Toeplitz algebra. It will be enough to show that the degree-0 part

$$(\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}} + \mathcal{K})^{(0)} = \pi^{-1}(\pi(\zeta(\mathcal{B}_\infty))) = (\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}})^{(0)} + \Gamma_0$$

is a C^* -algebra. But we have seen that $\zeta(fg) - \zeta(f)\zeta(g)$ is in Γ_0 for all $f, g \in \mathcal{B}_\infty$, so the map $\mathcal{B}_\infty \ni f \rightarrow \pi(\zeta(f))$ is a $*$ -homomorphism into Γ_b/Γ_0 whose image is thus a C^* -algebra and whose preimage in Γ_b is just $(\mathcal{A}_{\mathfrak{H}}^* \mathcal{A}_{\mathfrak{H}} + \mathcal{K})^{(0)}$. \square

Recall that it is the normally ordered elements of $\mathcal{T}_{\mathfrak{H}}^{(0)}$ which are constant under the inductive system $\iota_{\bullet, \bullet}$. The partial inverse $\varsigma^{(m)} : \varsigma^{(m)}(\mathcal{B}_m) \rightarrow \mathcal{B}_m$ of the covariant symbol map $\varsigma^{(m)}$ gives a normally ordered “quantization” of \mathcal{B}_∞ , and the image of $\prod_{m \in \mathbb{N}_0} \varsigma^{(m)}$ is the normally ordered part of the Toeplitz core, which is *all* of $\mathcal{T}_{\mathfrak{H}}^{(0)}$. In contrast, the projective limit \mathcal{B}^∞ contains (as an operator space) only the anti-normally ordered elements in $\mathcal{T}_{\mathfrak{H}}^{(0)}$, so “Toeplitz quantization” gives the anti-normal ordering. Lemma 2.15 shows that $\pi^{-1}(\pi(\zeta(\mathcal{B}_\infty))) = \zeta(\mathcal{B}_\infty) + \Gamma_0$ nevertheless gives all of $\mathcal{T}_{\mathfrak{H}}^{(0)}$.

5.8 Strict quantization

Some authors ([Haw3], [Rie1], [Sain]) do not require commutativity of the “classical limit algebra” in an axiomatic approach to “strict quantization”. Adapting such a definition, we can show that what we have done here is a strict quantization.

Let ${}^0\mathcal{B}_\infty$ denote the $*$ -algebra generated by the $\varsigma^{(m)}(A)$ ’s for all $A \in \mathcal{B}(\mathfrak{H}_m)$ and all $m \in \mathbb{N}_0$; thus ${}^0\mathcal{B}_\infty$ is a dense $*$ -subalgebra (the “algebraic part”) of the inductive-limit C^* -algebra \mathcal{B}_∞ .

Definition 5.25. The **Berezin product** on ${}^0\mathcal{B}_\infty$ is defined for all $f, g \in {}^0\mathcal{B}_\infty$ by

$$f \star^{(m)} g := \varsigma^{(m)}(\varsigma_{(m)}(f)\varsigma_{(m)}(g)),$$

where $\varsigma_{(m)} : \varsigma^{(m)}(\mathcal{B}_m) \rightarrow \mathcal{B}_m$ denotes the partial inverse of $\varsigma^{(m)}$. The **Poisson bracket** on ${}^0\mathcal{B}_\infty$ is defined by (cf. [Hawk1, §D.1])

$$\{f, g\} := \lim_{m \rightarrow \infty} \frac{m}{\sqrt{-1}} (f \stackrel{(m)}{\star} g - g \stackrel{(m)}{\star} f). \quad (5.21)$$

Of course we do not expect (5.21) to be a Poisson bracket in the ordinary sense, and $\{\cdot, \cdot\}$ is not likely to be interesting unless \mathfrak{H}_\bullet is commutative.

Corollary 5.26. *The sequence $(\mathcal{B}_\bullet, \zeta^{(\bullet)}) = (\mathcal{B}_m, \zeta^{(m)})_{m \in \mathbb{N}_0}$ is a strict quantization of ${}^0\mathcal{B}_\infty$ in the sense that each $\zeta^{(m)}$ is surjective and, for all $f, g \in {}^0\mathcal{B}_\infty$,*

$$\lim_{m \rightarrow \infty} \|\zeta^{(m)}(fg) - \zeta^{(m)}(f)\zeta^{(m)}(g)\| = 0, \quad (5.22)$$

$$\lim_{m \rightarrow \infty} \|\zeta^{(m)}(f)\| = \|f\|, \quad (5.23)$$

$$\lim_{m \rightarrow \infty} \|m^{-1}[\zeta^{(m)}(f), \zeta^{(m)}(g)] - \{f, g\}\| = 0. \quad (5.24)$$

Proof. We have seen that $\pi \circ \zeta : \mathcal{B}_\infty \rightarrow \mathcal{T}_{\mathfrak{H}}^{(0)}/\Gamma_0$ is injective, so

$$\zeta(f)\zeta(g) - \zeta(fg) \in \Gamma_0,$$

which is equivalent to von Neumann's condition (5.22). Rieffel's condition (5.23) coincides with the definition of the norm on \mathcal{B}^∞ . Similarly, the Dirac condition (5.24) is tautology in view of our definition of $\{\cdot, \cdot\}$ in (5.21). \square

We have seen that a subproduct system \mathfrak{H}_\bullet comes with a sequence $\mathcal{B}_\bullet = (\mathcal{B}_m)_{m \in \mathbb{N}_0}$ of finite-dimensional algebras $\mathcal{B}_m := \mathcal{B}(\mathfrak{H}_m)$, to which we can add $\mathcal{B}_\infty \cong \mathcal{O}_{\mathfrak{H}}^{(0)}$, and two sequences $\varsigma^{(\bullet)} = (\varsigma^{(m)})_{m \in \mathbb{N}_0}$ and $\zeta^{(\bullet)} = (\zeta^{(m)})_{m \in \mathbb{N}_0}$ of positive unital maps

$$\varsigma^{(m)} : \mathcal{B}_m \rightarrow \mathcal{B}_\infty, \quad \zeta^{(m)} : \mathcal{B}_\infty \rightarrow \mathcal{B}_m$$

such that $\varsigma^{(m)} \circ \zeta^{(m)}$ converges to the identity map on \mathcal{B}_∞ . As in [Sain, Prop. 2.2] we can associate to this data a continuous field of C^* -algebras, making explicit the assertion in Remark 4.5.

Corollary 5.27. *The C^* -algebra $\mathcal{T}_{\mathfrak{H}}^{(0)}$ can be identified with the space of continuous sections of the continuous field $\mathbb{N}_0 \cup \{\infty\} \ni m \rightarrow \mathcal{B}_m$, i.e.*

$$\mathcal{T}_{\mathfrak{H}}^{(0)} \cong \left\{ (X_m)_{m \in \mathbb{N}_0 \cup \{\infty\}} \in \prod_{m \in \mathbb{N}_0 \cup \{\infty\}} \mathcal{B}_m \mid X_\infty = \lim_{m \rightarrow \infty} \varsigma^{(m)}(X_m) \right\}.$$

Proof. We have seen that $\varsigma : \zeta(\mathcal{B}_\infty) \rightarrow \mathcal{B}_\infty$ can be obtained as

$$\varsigma(X) = \lim_{m \rightarrow \infty} \varsigma^{(m)}(Xp_m).$$

Thus, the C^* -algebra of continuous sections of \mathcal{B}_\bullet consists of the image of $\zeta(\mathcal{B}_\infty)$ under ς together with the sequences $(X_m)_{m \in \mathbb{N}_0 \cup \{\infty\}}$ such that $X_\infty = 0$. Hence the result follows from the facts that $\zeta(\mathcal{B}_\infty) + \Gamma_0 = \mathcal{T}_{\mathfrak{H}}^{(0)}$ and that ς is an isomorphism. \square

5.9 $\mathcal{O}_{\mathfrak{H}}$ assembled from projective limits

We now define modules over the projective limit \mathcal{B}^∞ .

Recall that we defined in §4.3 an inductive system $\iota_{m,l}^{(k)} : \mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+k}) \rightarrow \mathcal{B}(\mathfrak{H}_l, \mathfrak{H}_{l+k})$. Define the adjoint $j_{l,m}^{(k)} : \mathcal{B}(\mathfrak{H}_{l+k}, \mathfrak{H}_l) \rightarrow \mathcal{B}(\mathfrak{H}_{m+k}, \mathfrak{H}_m)$ of $\iota_{m,l}^{(k)}$ by the property that

$$\phi_m(j_{l,m}^{(k)}(X)Y) = \phi_l(X\iota_{m,l}^{(k)}(Y))$$

for all $X \in \mathcal{B}(\mathfrak{H}_{l+k}, \mathfrak{H}_l)$ and all $Y \in \mathcal{B}(\mathfrak{H}_m, \mathfrak{H}_{m+k})$. We deduce that

$$j_{l,m}^{(k)}(X) = \frac{\text{Tr}(Q_m)}{\text{Tr}(Q_l)} \sum_{|\mathbf{r}|=l-m} (Q^{\otimes m})_{\mathbf{r},\mathbf{r}} R_{\mathbf{r}}^* X R_{\mathbf{r}}|_{\mathfrak{H}_m}, \quad \forall X \in \mathcal{B}(\mathfrak{H}_{l+k}, \mathfrak{H}_l),$$

and that the operators on $\mathfrak{H}_{\mathbb{N}}$ which are constant with respect to the system $j_{\bullet,\bullet}^{(k)}$ are precisely those of the form $\check{\zeta}_k(f) = (\check{\zeta}_k^{(l)}(f))_{l \in \mathbb{N}_0}$ for some $f \in \mathcal{B}_\infty$, where

$$\check{\zeta}_k^{(l)}(f) = \text{Tr}(Q_l) \sum_{|\mathbf{j}|=l, |\mathbf{k}|=l+k} (Q^{\otimes m})_{\mathbf{j},\mathbf{j}}^{-1} \omega_Q(Z_{\mathbf{j}} Z_{\mathbf{k}}^* f) S_{\mathbf{j}} S_{\mathbf{k}}^*|_{\mathfrak{H}_l}.$$

Define $\mathcal{E}_{(k)}$ to be the set of $\check{\zeta}_k(f)$'s for all $f \in \mathcal{B}_\infty$. Then $\mathcal{E}_{(k)}$ is an operator system.

Let $\Gamma_0^{(k)}$ be the vector space of sequences of operators in $\mathcal{B}(\mathfrak{H}_{l+k}, \mathfrak{H}_l)$ which converge to zero as $l \rightarrow \infty$.

Proposition 5.28. *The vector space $\mathcal{E}_{(k)} + \Gamma_0^{(k)}$ coincides with the subspace $\mathcal{T}_{\mathfrak{H}}^{(-k)}$ of the Toeplitz algebra. Hence $\mathcal{E}_{(k)}$ is a module over the C^* -algebra \mathcal{B}^∞ , the module action taken modulo $\Gamma_0^{(k)}$. In fact, $\mathcal{E}_{(k)}$ is isomorphic as a Hilbert bimodule to $\mathcal{E}^{(-k)} \cong \mathcal{O}_{\mathfrak{H}}^{(-k)}$.*

Proof. The first statements are proven in the same way as for $k = 0$. For the last assertion, note that the algebras of compact module operators $\mathcal{K}_{\mathcal{B}_\infty}(\mathcal{E}^{(-k)})$ and $\mathcal{K}_{\mathcal{B}_\infty}(\mathcal{E}_{(k)})$ are isomorphic, namely to $\mathcal{B}_\infty \cong \mathcal{B}^\infty$. Hence the modules $\mathcal{E}_{(k)}$ and $\mathcal{E}^{(-k)}$ are isomorphic [Frank1]. \square

5.10 Commutative case and the Arveson conjecture

In [Vas1], [Vas3] it was shown that the Cuntz–Pimsner algebra $\mathcal{O}_{\mathcal{E}}$ (defined in [Pims1]) of the $C(M)$ -Hilbert bimodule \mathcal{E} of continuous sections of a line bundle $L \rightarrow M$ is isomorphic to the C^* -algebra $C(\mathbb{S}_L)$ of continuous functions on the total space of the circle bundle \mathbb{S}_L associated to L^* . Recall that in the definition of $\mathcal{O}_{\mathcal{E}}$ (which is Pimsner's original one) the tensor products are taken over the coefficient algebra $C(M)$. We shall now see that, in the case (M, L) is a projectively induced quantization, we can also obtain $C(\mathbb{S}_M) := C(\mathbb{S}_L)$ as the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ of the associated subproduct system \mathfrak{H}_\bullet . The Hilbert space \mathfrak{H}_m is the “holomorphic part” of the module $\mathcal{E}^{\otimes m}$ (where $\otimes := \otimes_{C(M)}$).

Proposition 5.29. *Let (M, L) be a polarized (not necessarily smooth) variety and let \mathfrak{H}_\bullet be a projectively induced quantization of (M, L) (the definition still makes sense in the non-smooth case), and endow M with the complex (Hausdorff) topology [Serre1, §2]. Then*

$$C(\mathbb{S}_M) \cong \mathcal{O}_{\mathfrak{H}}, \tag{5.25}$$

and for all $k \in \mathbb{Z}$,

$$\Gamma(M, L^{\otimes k}) \cong \mathcal{O}_{\mathfrak{H}}^{(k)}$$

as a Hilbert $C(M)$ -bimodule. In particular, $k = 0$ gives $C(M)$ as an inductive limit.

Proof. As in the general case, $\mathcal{O}_{\mathfrak{H}}$ is built up from Hilbert modules over $\mathcal{O}_{\mathfrak{H}}^{(0)}$ and the latter is generated by the images of the covariant symbol maps $\varsigma^{(m)}$. An argument given in [CGR1, §4] shows that the $\varsigma^{(m)}(A)$'s (for all $m \in \mathbb{N}_0$ and all $A \in \mathcal{B}_m$) separate points. The Stone–Weierstrass theorem gives that they form a dense subalgebra of $C(M)$. The inclusion of \mathfrak{H}_m into \mathfrak{H}_{m+1} coincide by construction with our $\iota_{m,m+1}$, and the supremum norm on $C(M)$ is seen to coincide with the norm on the inductive limit \mathcal{B}_∞ . Therefore, $C(M) \cong \mathcal{B}_\infty$. The result now follows from Theorem 4.12 and the well-know decomposition of $C(\mathbb{S}_M)$ into the $\Gamma(M, L^{\otimes m})$'s (see §3.3). \square

Proposition 5.29 was inspired [Haw2], where the focus lied on the Toeplitz operators. When M is smooth we see from the proof of [Haw2, Lemma 4.2] that the limit state on $\mathcal{B}_\infty = C(M)$ is faithful and hence $C(M)$ is also isomorphic to the subset $\zeta(\mathcal{B}_\infty)$ of the projective limit \mathcal{B}^∞ with the multiplication taken modulo compacts. For non-smooth M we do not know if the limit state on \mathcal{B}_∞ is faithful.

Note that it is not so obvious that we could recover M completely (as a topological space) from \mathfrak{H}_\bullet because the homogeneous coordinate ring $\bigoplus_{m \in \mathbb{N}_0} H^0(M; L^{\otimes m})$ of $M \subset \mathbb{P}[\mathfrak{H}^*]$ is not a ring of functions on M . The choice of basis on $H^0(M; L^{\otimes m})$ corresponds to a choice of algebra structure on the ring $C(M)$ and the inner product on $H^0(M; L^{\otimes m})$ to a choice of C^* -algebra structure on $C(M)$, but all possible C^* -algebra structures obtain in this way are isomorphic.

In the following we use the terminology from Lemma 3.8.

Corollary 5.30. *Let (M, L) be a polarized manifold and let \mathfrak{H}_\bullet be a projectively induced quantization of (M, L) . Then the limit state on $C(M) := \mathcal{B}_\infty$ coincides with the unique volume form on M which balances the inner product on \mathfrak{H} . The Fubini–Study metric $\text{FS}(\langle \cdot | \cdot \rangle)$ on L associated with the inner product on \mathfrak{H} coincides with the $*$ -operation which defines the C^* -algebra $\mathcal{O}_{\mathfrak{H}}$, and is thus equal to the inductive limit of the Hermitian pairings*

$$\mathcal{B}(\mathfrak{H}_{m+1}, \mathfrak{H}_m) \times \mathcal{B}(\mathfrak{H}_{m+1}, \mathfrak{H}_m) \rightarrow \mathcal{B}(\mathfrak{H}_m), \quad (A, B) \rightarrow A^*B.$$

Consequently, (M, L) is balanced if and only if the limit state coincides with the normalized Fubini–Study volume form restricted to $M \subset \mathbb{P}[\mathfrak{H}^]$.*

Proof. We just have to recall (cf. Lemma 3.8) that the generators Z_1, \dots, Z_n of $\mathcal{O}_{\mathfrak{H}}$ satisfy $\sum_{k=1}^n Z_k Z_k^* = 1$ and that the limit state $\omega_{p_1} : C(M) \rightarrow \mathbb{C}$ satisfies $\omega_{p_1}(Z_j^* Z_k) = \delta_{j,k}/n$. \square

As mentioned, the existence of the $\langle \cdot | \cdot \rangle$ -balancing volume form ω_{p_1} is related to the Calabi–Yau theorem, and it was previously known that ω_{p_1} can be obtained using finite-dimensional approximation [CaKe1], [Don3, §2.2]. This procedure comes out automatically in a slightly different guise in our approach using inductive limits; the limit state on \mathcal{B}_∞ is a generalization of the volume form associated with the Calabi–Yau metric.

By [Zeld1], the Hermitian metric h on L used to define the inner product on \mathfrak{H} is recovered the C^∞ -topology as the $m \rightarrow \infty$ limit of the pullbacks of the Fubini–Study metrics via the Kodaira embeddings. In a rather different fashion, Corollary 5.30 gives the Fubini–Study Hermitian metric $\text{FS}(\langle \cdot | \cdot \rangle)$ on L as a limit of the inner products of the \mathfrak{H}_m 's. That is, we obtain a Hermitian metric on L from a sequence of matrix-valued *inner products* on Hilbert modules over finite-dimensional matrix algebra. These two approximation results are rather similar though since the space of pullback metrics with respect to a Kodaira embeddings for $H^0(M; L^m)$ identifies with the space of inner products on $H^0(M; L^m)$.

As we have seen (recall Proposition 3.6), from the version of Berezin quantization with prequantum condition one obtains a strict quantization of $C(M)$. With projectively induced quantization we obtain from Corollary 5.26 a strict quantization of $C(M)$, and we do not require M to be smooth.

Corollary 5.31. *For any projective variety M , the sequence $(\mathcal{B}_m, \zeta^{(m)})_{m \in \mathbb{N}_0}$ is a strict quantization of the dense $*$ -subalgebra of $C(M)$ generated by the $\varsigma^{(m)}(\mathcal{B}_m)$'s.*

Finally we arrive at one of the most striking applications of our results, namely to the Arveson conjecture (see Remark 2.17). The validity of this conjecture has been proven by other means in many cases [DoWa2], [EnEs1].

Corollary 5.32. *Arveson's conjecture holds for all homogeneous ideals $\mathcal{I} \subset \mathbb{C}[z_1, \dots, z_n]$, i.e. the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ of the subproduct system \mathfrak{H}_\bullet associated to \mathcal{I} (as in Lemma 2.9) is commutative.*

In fact, Corollary 5.32 follows directly from Lemma 2.15 if we use [KeSh1, Prop. 4.14].

6 Application to compact matrix quantum groups

6.1 Compact matrix quantum groups

For the theory of compact quantum groups we refer to [KLS], [MaVD], [Timm1]. We shall restrict attention to compact *matrix* quantum groups, defined as follows.

Definition 6.1 ([Wang3], [Wor1]). A **compact matrix quantum group** \mathbb{G} is defined by a C^* -algebra $C(\mathbb{G})$ generated by the entries $u_{j,k}$ of a single unitary matrix $u \in M_n(\mathbb{C}) \otimes C(\mathbb{G})$ (for some $n \in \mathbb{N}$) such that the map $\Delta : C(\mathbb{G}) \rightarrow C(\mathbb{G}) \otimes C(\mathbb{G})$ defined by

$$\Delta(u_{j,k}) := \sum_{r=1}^n u_{j,r} \otimes u_{r,k}$$

is a $*$ -homomorphism, and such that the transpose u^t is invertible.

We refer to the generating matrix u as the **defining representation** of the “group” \mathbb{G} .

The **Haar state** on $C(\mathbb{G})$ (or the **Haar measure** on \mathbb{G}) is the unique state on $C(\mathbb{G})$ which is **left \mathbb{G} -invariant**, in the sense that

$$(\text{id} \otimes h) \circ \Delta(f) = h(f)\mathbf{1}.$$

The Haar state is always faithful on the $*$ -algebra generated by the $u_{j,k}$'s but not necessarily so on the norm closure $C(\mathbb{G})$. There is a canonical construction of a “reduced version” of \mathbb{G} , which is a compact quantum group with faithful Haar state [BMT1, §2] and has the same dense Hopf $*$ -algebra. We shall always work with the reduced version or, what amounts to the same thing, assume that h is faithful on all of $C(\mathbb{G})$. Then h is a KMS state [Wor4].

6.1.1 Representations and actions

Definition 6.2. A **representation** of \mathbb{G} is a **corepresentation** of $C(\mathbb{G})$, i.e. an invertible element $v \in \mathcal{B}(\mathfrak{H}_v) \otimes C(\mathbb{G})$, for some Hilbert space \mathfrak{H}_v , satisfying (in leg-numbering notation)

$$(\text{id} \otimes \Delta)(v) = v_{13}v_{23}$$

as elements of $\mathcal{B}(\mathfrak{H}_v) \otimes \mathcal{B}(\mathfrak{H}_v) \otimes C(\mathbb{G})$. A representation v is **irreducible** if the set

$$\text{Hom}_{\mathbb{G}}(v, v) := \{T \in \mathcal{B}(\mathfrak{H}_v) \mid (T \otimes \mathbf{1})v = v(T \otimes \mathbf{1})\}$$

is trivial.

Definition 6.3. Two representations $v \in \mathcal{B}(\mathfrak{H}_v) \otimes C(\mathbb{G})$ and $w \in \mathcal{B}(\mathfrak{H}_w) \otimes C(\mathbb{G})$ are **equivalent**, denoted $v \simeq w$, if there is a unitary $U : \mathfrak{H}_v \rightarrow \mathfrak{H}_w$ such that

$$(U \otimes \mathbf{1})v = w(U \otimes \mathbf{1})$$

(in particular, this requires $\dim \mathfrak{H}_v = \dim \mathfrak{H}_w$). We denote by $\text{Irrep } \mathbb{G}$ the (countable) set of equivalence classes of irreducible representations of \mathbb{G} . We choose a representative $u^{(\lambda)} \in \mathcal{B}(\mathfrak{H}_\lambda) \otimes C(\mathbb{G})$ for each $\lambda \in \hat{\mathbb{G}}$.

Definition 6.4. The **tensor product** of two representations $u \in \mathcal{B}(\mathfrak{H}) \otimes C(\mathbb{G})$ and $v \in \mathcal{B}(\mathfrak{K}) \otimes C(\mathbb{G})$ is the representation

$$u \otimes v := u_{13}v_{23} \in \mathcal{B}(\mathfrak{H} \otimes \mathfrak{K}) \otimes C(\mathbb{G}).$$

In particular, $u^{\otimes m} = u_{1,m+1} \cdots u_{m,m+1}$ is the matrix whose entries in the product basis for $\mathfrak{H}^{\otimes m}$ is given by

$$u_{\mathbf{j}, \mathbf{k}} = u_{j_1, k_1} \cdots u_{j_m, k_m}.$$

Now let us explain the motivation for the invertible operator $Q \in \mathcal{B}(\mathfrak{H})$ that we incorporated in the Berezin quantization (recall §4.5).

It is a crucial consequence of the axioms of compact matrix quantum groups that for any finite-dimensional representation $v \in \mathcal{B}(\mathfrak{H}_v) \otimes C(\mathbb{G})$ of \mathbb{G} , one can find an invertible matrix $F_v \in \mathcal{B}(\mathfrak{H}_v)$ such that

$$\bar{v} := (F_v \otimes \mathbf{1})v^c(F_v^{-1} \otimes \mathbf{1}) \quad (6.1)$$

is unitary, where $v^c = (v^t)^*$ is the matrix whose coefficients are the adjoints of those of v . The equivalence class of \bar{v} is the **conjugate** of the equivalence class of v (we shall also say that \bar{v} is a conjugate of v). The matrix F_v in (6.1) is usually chosen such that $Q_v := F_v^* F_v$ satisfies

$$\mathrm{Tr}(Q_v^{-1}) = \mathrm{Tr}(Q_v) \equiv \dim_q(v),$$

and this quantity is the “quantum dimension” of v . We have $Q_{\bar{v}} = (Q_v^t)^{-1}$. We shall write $Q_\lambda := Q_{u(\lambda)}$ etc. for irreducibles $\lambda \in \mathrm{Irrep} \mathbb{G}$ and we denote by $\bar{\lambda}$ the conjugate of λ .

Every representation of \mathbb{G} decomposes completely into a direct sum of irreducibles. Hence, for each pair of irreps $\lambda, \mu \in \mathbb{G}$ there are integers $\mathrm{mult}(\nu, \lambda \otimes \mu) \in \mathbb{N}_0$ such that

$$u^{(\lambda)} \otimes u^{(\mu)} \simeq \bigoplus_{\nu \in \mathrm{Irrep} \mathbb{G}} \mathrm{mult}(\nu, \lambda \otimes \mu) u^{(\nu)}. \quad (6.2)$$

Definition 6.5. The equations (6.2) dictate the **fusion rules** of \mathbb{G} . The fusion rules are **commutative** if

$$\mathrm{mult}(\nu, \lambda \otimes \mu) = \mathrm{mult}(\lambda \otimes \mu, \nu), \quad \forall \lambda, \mu, \nu \in \mathrm{Irrep} \mathbb{G}.$$

Example 6.6. Compact groups $\mathbb{G} = G$ have commutative fusion rules. More generally, q -deformations of compact Lie groups have commutative fusion rules because the equivalence class of an irreducible representation is determined by the highest weight of the representation.

The quantum groups in the next two examples are introduced in Definition 6.10 below.

Example 6.7. For any F , the fusion rules of the quantum group $B_u(F)$ are identical to those of $\mathrm{SU}(2)$; in particular this is true for $\mathrm{SU}_q(2)$. These fusion rules in fact characterize the $B_u(F)$ ’s among compact quantum groups [Ban3, Théorème 2].

Example 6.8. The fusion rules of $A_u(Q)$ are far from commutative, see [Ban4].

Definition 6.9 ([Wang1, Def. 3.1]). A **left action** of a compact matrix quantum group \mathbb{G} on a C^* -algebra \mathcal{B} is a unital $*$ -homomorphism $\alpha : \mathcal{B} \rightarrow \mathcal{B} \otimes C(\mathbb{G})$ such that

- (i) $(\alpha \otimes \mathrm{id}) \circ \alpha = (\mathrm{id} \otimes \Delta) \circ \alpha$,
- (ii) $(\mathrm{id} \otimes \varepsilon) \circ \alpha = \mathrm{id}$, where ε is the counit on the dense Hopf- $*$ -subalgebra of $C(\mathbb{G})$, and
- (iii) there is a dense $*$ -subalgebra ${}^0\mathcal{B}$ of \mathcal{B} such that $\alpha({}^0\mathcal{B}) \subset {}^0\mathcal{B} \otimes C^\infty(\mathbb{G})$.

Similarly, a **right action** of \mathbb{G} on \mathcal{B} is a unital $*$ -homomorphism $\alpha : \mathcal{B} \rightarrow C(\mathbb{G}) \otimes \mathcal{B}$ satisfying the obvious analogues of the properties (i), (ii) and (iii).

If \mathcal{B} is a von Neumann algebra then we replace $C(\mathbb{G})$ by its weak closure $L^\infty(\mathbb{G})$ in the GNS representation of the Haar state, and only condition (i) is required in the definition of an action.

Every unitary representation $v \in \mathcal{B}(\mathfrak{H}) \otimes C(\mathbb{G})$ of \mathbb{G} induces a left action of \mathbb{G} on $\mathcal{B}(\mathfrak{H})$ given by

$$\mathrm{Ad}_v : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H}) \otimes L^\infty(\mathbb{G}), \quad \mathrm{Ad}_v(A) := v(A \otimes \mathbf{1})v^*. \quad (6.3)$$

6.1.2 Universal quantum groups

In the following, for a matrix u with entries in $C(\mathbb{G})$, we write u^c for the transpose of u^* , i.e. $(u^c)_{j,k} := u_{j,k}^*$ where $u_{j,k}^*$ is the adjoint of $u_{j,k}$ in $C(\mathbb{G})$.

Definition 6.10 ([Wang3], [Ban4, Déf. 1]). Let $F \in \text{GL}(n, C)$ be an invertible matrix and write $Q := F^*F$. The **universal unitary quantum group** $\mathbb{G} = A_u(Q)$ is the compact matrix quantum group \mathbb{G} whose algebra of continuous functions $C(\mathbb{G})$ is generated by the entries of a unitary $n \times n$ matrix u satisfying the relations making $(F \otimes \mathbf{1})u^c(F^{-1} \otimes \mathbf{1})$ a unitary matrix.

The **universal orthogonal quantum group** $\mathbb{G} = B_u(F)$ is the compact matrix quantum group whose algebra $C(\mathbb{G})$ is the quotient of that of $A_u(Q)$ by the relation $u = (F \otimes \mathbf{1})u^c(F^{-1} \otimes \mathbf{1})$.

The prototype example of a $B_u(F)$ is the quantum $\text{SU}_q(2)$ group $\mathbb{G} := \text{SU}_q(2)$. In general, $B_u(F)$ is some kind of higher-dimensional quantum $\text{SU}(2)$ group which has no classical counterpart.

Suppose that \mathbb{H} and \mathbb{G} are compact matrix quantum groups such that $C(\mathbb{H})$ is a quotient of $C(\mathbb{G})$. If the quotient map $\pi : C(\mathbb{G}) \rightarrow C(\mathbb{H})$ fulfills $(\pi \otimes \pi) \circ \Delta_{\mathbb{G}} = \Delta_{\mathbb{H}} \circ \pi$, i.e. if π intertwines the comultiplication of \mathbb{G} with that of \mathbb{H} , then \mathbb{H} is a **quantum subgroup** \mathbb{G} . We have seen that $B_u(F)$ is a quantum subgroup of $A_u(Q)$ when $F^*F = Q$.

The name “universal” is motivated by the following fact, which we should anticipate from (6.1).

Lemma 6.11 ([VaDW]). *Any compact matrix quantum group \mathbb{G} is a quantum subgroup of $A_u(Q)$ for some Q . If \mathbb{G} in addition has a self-conjugate defining representation, then $C(\mathbb{G})$ is a quantum subgroup of some $B_u(F)$. We write $\mathbb{G} \subset A_u(Q)$ and $\mathbb{G} \subset B_u(F) \subset A_u(Q)$ for these cases respectively.*

Let u be the fundamental representation of $\mathbb{G} \subset A_u(Q)$, with $Q \in \text{GL}(n, \mathbb{C})$. Then the elements $z_1 := u_{k,1}, \dots, z_n := u_{k,n}$ of the first row of u satisfy the **Q -sphere relations**

$$(Q^{-1})_{1,1} \mathbf{1} = \sum_{r,s=1}^n (Q^{-1})_{r,s} z_r^* z_s, \quad \mathbf{1} = \sum_{s=1}^n z_s z_s^*. \quad (6.4)$$

6.1.3 The dual discrete quantum group

Let \mathbb{G} be a compact matrix quantum group such that the GNS representation $C(\mathbb{G}) \rightarrow \mathcal{B}(L^2(\mathbb{G}))$ of the Haar state is faithful. We shall identify $C(\mathbb{G})$ with its image in $\mathcal{B}(L^2(\mathbb{G}))$ and denote by $L^\infty(\mathbb{G})$ the von Neumann algebra generated by $C(\mathbb{G})$ in $\mathcal{B}(L^2(\mathbb{G}))$.

In perfect analogy to the case of ordinary compact groups, the C^* -algebra $C(\mathbb{G})$ has a **Peter–Weyl decomposition**

$$C(\mathbb{G}) = \bigoplus_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda)^*, \quad (6.5)$$

and the completion $L^2(\mathbb{G})$ of $C(\mathbb{G})$ in the inner product defined by the Haar state then allows for a similar decomposition. The comultiplication Δ on $C(\mathbb{G}) \subset \mathcal{B}(L^2(\mathbb{G}))$ takes the form

$$\Delta(f) = W(f \otimes \mathbf{1})W^*, \quad \forall f \in C(\mathbb{G})$$

for a unitary operator W on $L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ referred to as the **multiplicative unitary** of \mathbb{G} . The **dual** of \mathbb{G} is then defined via the (multiplier) Hopf C^* -algebra

$$c_0(\hat{\mathbb{G}}) := \bigoplus_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda), \quad (6.6)$$

with the comultiplication given by

$$\hat{\Delta}(X) := W^*(\mathbf{1} \otimes X)W, \quad \forall X \in c_0(\hat{\mathbb{G}}).$$

We denote by p_λ the identity in $\mathcal{B}(\mathfrak{H}_\lambda)$, regarded as an element of $c_0(\hat{\mathbb{G}})$. Then the counit $\hat{\varepsilon}$ on $c_0(\hat{\mathbb{G}})$ is characterized by (cf. (2.4))

$$\hat{\varepsilon}(X)p_0 = Xp_0, \quad \forall X \in c_0(\hat{\mathbb{G}}),$$

where $0 \in \text{Irrep } \mathbb{G}$ is the trivial representation. Every irreducible representation $u^{(\lambda)}$ of \mathbb{G} is obtained from W by means of

$$u^{(\lambda)} = W(p_\lambda \otimes \mathbf{1}).$$

The object $\hat{\mathbb{G}}$ is referred to as a **discrete quantum group**. In the general theory of “locally compact quantum groups”, there is a canonical dual quantum group associated also to $\hat{\mathbb{G}}$, and this quantum group is precisely \mathbb{G} . In particular, the dual of an ordinary compact group G is a discrete quantum group, which is an honest group only if G is abelian.

The C^* -algebra $c_0(\hat{\mathbb{G}})$ is contained in the C^* -algebra \mathcal{K} of compact operators on $L^2(\mathbb{G})$. The multiplier algebra of $c_0(\hat{\mathbb{G}})$ can be identified with the C^* -direct product

$$c_b(\hat{\mathbb{G}}) := \prod_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda),$$

and the comultiplication $\hat{\Delta}$ is a map from $c_0(\hat{\mathbb{G}})$ into $c_b(\hat{\mathbb{G}}) \otimes c_b(\hat{\mathbb{G}})$. Continuing the analogy with the theory of honest groups, we shall denote by

$$c_c(\hat{\mathbb{G}}) := \bigoplus_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda)$$

the algebraic sum, and we denote the weak closure of $c_0(\hat{\mathbb{G}})$ in $\mathcal{B}(L^2(\mathbb{G}))$ by

$$\mathcal{R}(\mathbb{G}) = \ell^\infty(\hat{\mathbb{G}}) := \ell^\infty - \prod_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda).$$

This “group-von Neumann algebra” $\mathcal{R}(\mathbb{G})$ is contained in the dual $C(\mathbb{G})^*$ of $C(\mathbb{G})$.

Finally, the algebra of operators on $L^2(\mathbb{G})$ affiliated with $\mathcal{R}(\mathbb{G})$ can be identified with the product $\hat{\prod}_{\lambda \in \text{Irrep } \mathbb{G}} \mathcal{B}(\mathfrak{H}_\lambda)$, containing all (not necessarily bounded) sequences of elements in the $\mathcal{B}(\mathfrak{H}_\lambda)$ ’s. Of particular importance is the operator $Q_{\mathbb{G}} := (Q_\lambda)_{\lambda \in \text{Irrep } \mathbb{G}}$, where Q_λ is as in §6.1.1.

From (6.6) we see that the irreducible representations of the C^* -algebra $c_0(\hat{\mathbb{G}})$ (also referred to as the irreducible “corepresentations” of $\hat{\mathbb{G}}$) are parameterized by $\lambda \in \text{Irrep } \mathbb{G}$. In fact, if $u^{(\lambda)} \in \mathcal{B}(\mathfrak{H}_\lambda) \otimes C(\mathbb{G})$ is the irreducible representation of \mathbb{G} with label λ (as in §6.1.1) then

$$\pi_\lambda(X) := (\text{id} \otimes X)(u^{(\lambda)}) \tag{6.7}$$

is the corresponding irreducible corepresentation of $\hat{\mathbb{G}}$, where $X \in c_0(\hat{\mathbb{G}})$ is regarded as a functional on $C(\mathbb{G})$. In general, if v is a unitary representation of \mathbb{G} then (6.7) defines a representation π_v of $\hat{\mathbb{G}}$ by substituting $u^{(\lambda)}$ with v . Then the commutant of $\pi_v(c_0(\hat{\mathbb{G}}))$ in $\mathcal{B}(\mathfrak{H}_v)$ equals

$$\pi_v(c_0(\hat{\mathbb{G}}))' = \mathcal{B}(\mathfrak{H}_v)^{\mathbb{G}},$$

the fixed-point subalgebra under the \mathbb{G} -action Ad_v (recall (6.3)).

The same equation (6.7) represents any (possibly unbounded) operator X affiliated to $\mathcal{R}(\mathbb{G})$ on $\mathcal{B}(\mathfrak{H}_\lambda)$. For each finite-dimensional representation v of \mathbb{G} ,

$$\pi_v(Q_{\mathbb{G}}) = Q_v^t,$$

where Q_v is as in §6.1.1 and Q_v^t denotes its transpose. In particular, Q_v^t commutes with the projection onto any irreducible subrepresentation of v . It is well known that $\hat{\Delta}(Q_{\mathbb{G}}) = Q_{\mathbb{G}} \otimes Q_{\mathbb{G}}$, and it gives

$$Q_{v \otimes w} = Q_v \otimes Q_w$$

for all finite-dimensional representations v and w .

6.2 First-row and first-column algebras

Throughout this section, \mathbb{G} is a compact matrix quantum group. We denote by (u, \mathfrak{H}) the defining representation of \mathbb{G} . Thus the C^* -algebra $C(\mathbb{G})$ is generated by the matrix coefficients of the unitary matrix $u \in \mathcal{B}(\mathfrak{H}) \otimes C(\mathbb{G})$. Set $n := \dim(\mathfrak{H})$ and fix an orthonormal basis e_1, \dots, e_n of \mathfrak{H} so that $\mathfrak{H} \cong \mathbb{C}^n$, and let $u_{j,k}$ be the matrix coefficients of u in this basis.

Definition 6.12. The **first-row algebra** of \mathbb{G} is the C^* -algebra $C(\mathbb{S}_{\mathbb{G}})$ generated by the first row $z_1 := u_{1,1}, \dots, z_n := u_{1,n}$. This defines the quantum homogeneous space $\mathbb{S}_{\mathbb{G}}$.

There is a \mathbb{Z} -grading on $C(\mathbb{S}_{\mathbb{G}})$ obtained by letting the z_k 's have degree 1 while their adjoints are given degree -1 . We write the decomposition into spectral subspaces for the corresponding $U(1)$ -action as

$$C(\mathbb{S}_{\mathbb{G}}) = \overline{\bigoplus_{k \in \mathbb{Z}} C(\mathbb{S}_{\mathbb{G}})^{(k)}}^{\|\cdot\|}.$$

Definition 6.13. The quantum homogeneous space \mathbb{G}/\mathbb{K} is defined as the noncommutative manifold corresponding to the C^* -subalgebra of fixed points in $C(\mathbb{S}_{\mathbb{G}})$ for the $U(1)$ -action:

$$C(\mathbb{G}/\mathbb{K}) := C(\mathbb{S}_{\mathbb{G}})^{(0)}.$$

It is clear that $C(\mathbb{G}/\mathbb{K})$ is generated by the n^2 elements $\{z_j^* z_k\}_{j,k=1}^n$ (but this is obviously not a minimal set of generators).

Example 6.14. If $\mathbb{G} = G$ is a compact semisimple Lie group then \mathbb{G}/\mathbb{K} is a coadjoint orbit and $\mathbb{S}_{\mathbb{G}}$ is a principal $U(1)$ -bundle over \mathbb{G}/\mathbb{K} . Indeed, let $(U_{-1}, \mathfrak{H}_{-1})$ be the irreducible unitary representation of G with highest weight $(-1, 0, \dots, 0)$ and let K be the stabilizer of the complex line spanned by the highest-weight vector ξ_{-1} . The action U_{-1} of G on \mathfrak{H}_{-1} is unitary, and hence Hamiltonian for the symplectic form given by the imaginary part of the inner product on \mathfrak{H}_{-1} . The orbit $U_{-1}(G) \cdot [\xi_{-1}]$ of the line spanned by ξ_{-1} is then a Hamiltonian G -homogeneous space, and a characterization of coadjoint orbits shows that $G/K \cong U_{-1}(G) \cdot [\xi_{-1}]$ is a coadjoint orbit.

The hyperplane bundle over the projectivization $\mathbb{P}[\mathfrak{H}_{-1}]$ restricts to a holomorphic line bundle L over G/K and we let \mathbb{S}_G be the total space of the principal $U(1)$ -bundle over G/K associated with L^* (cf. §3.3). A basis for the space of holomorphic sections of L generates $C(\mathbb{S}_G)$ and, after fixing a basis e_1^*, \dots, e_n^* for $\mathfrak{H}_{-1} \cong \mathbb{C}^n$, such a basis is provided by the coordinate functions z_1, \dots, z_n of \mathbb{CP}^{n-1} restricted to G/K . Choosing the basis such that $e_1^* \propto \xi_{-1}$ we get $\mathbb{G}/\mathbb{K} = G/K$ and $\mathbb{S}_{\mathbb{G}} = \mathbb{S}_G$.

For example, if $G = \mathrm{SU}(n)$ then $K = U(1) \times \mathrm{SU}(n-1)$ and $\mathbb{G}/\mathbb{K} = G/K$ is complex projective n -space \mathbb{CP}^{n-1} , whereas $\mathbb{S}_{\mathrm{SU}(n)} = \mathbb{S}^{2n-1}$ is the unit sphere in \mathbb{C}^n .

Example 6.15. The preceding discussion carries over to q -deformations of G , and we get that \mathbb{G}/\mathbb{K} is a quantum flag manifold. For $\mathbb{G} = \mathrm{SU}_q(n)$ we obtain quantum projective n -space $\mathbb{G}/\mathbb{K} = \mathbb{CP}_q^{n-1}$, and $\mathbb{S}_{\mathbb{G}}$ is the q -deformed $(2n-1)$ -sphere \mathbb{S}_q^{2n-1} (cf. [DDL1]).

Example 6.16 ([BaGo1, Thm. 3.3]). Let $\mathbb{G} := \mathrm{O}^*(n)$ be the half-liberated orthogonal group. Then $C(\mathbb{G}/\mathbb{K})$ is in fact commutative and \mathbb{G}/\mathbb{K} is just the ordinary complex projective n -space \mathbb{CP}^{n-1} ,

$$C(\mathbb{G}/\mathbb{K}) \cong C(\mathbb{CP}^{n-1}).$$

The spectral subspaces $C(\mathbb{S}_{\mathbb{G}})^{(k)}$ for the $U(1)$ action on $C(\mathbb{S}_{\mathbb{G}})$ are Hilbert bimodules over the fixed-point subalgebra $C(\mathbb{G}/\mathbb{K})$, where the multiplication in the ambient algebra $C(\mathbb{S}_{\mathbb{G}})$ defines left and right $C(\mathbb{G}/\mathbb{K})$ -valued inner products

$$\langle \xi | \eta \rangle_{\text{right}} := \xi^* \eta, \quad \langle \xi | \eta \rangle_{\text{left}} := \xi \eta^* \tag{6.8}$$

between elements ξ and η in $C(\mathbb{S}_{\mathbb{G}})^{(k)}$.

Remark 6.17 (Row vs column). We can also consider the C^* -algebra generated by the first column elements $w_j := u_{j,1}$ of u (the “first-column algebra”). Everything proven about the first-row algebra $C(\mathbb{S}_{\mathbb{G}})$ in this paper has a version where one instead uses the generators of the first-column algebra.

Lemma 6.18. *The first-row algebra $C(\mathbb{S}_{\mathbb{G}})$ carries an ergodic action of \mathbb{G} which contains every irreducible representation of \mathbb{G} with multiplicity one.*

Proof. We define a left action $C(\mathbb{S}_{\mathbb{G}}) \rightarrow C(\mathbb{S}_{\mathbb{G}}) \otimes C(\mathbb{G})$ by restriction of the comultiplication. The Peter–Weyl decomposition (6.5) of $C(\mathbb{G})$ gives the decomposition

$$C(\mathbb{S}_{\mathbb{G}}) \cong \bigoplus_{\lambda \in \text{Irrep } \mathbb{G}} \mathfrak{H}_{\lambda},$$

and the comultiplication restricts to the irreducible \mathbb{G} -representation $u^{(\lambda)}$ on each \mathfrak{H}_{λ} . \square

In view of Lemma 6.18 and the above examples, the quantum homogeneous \mathbb{G} -space \mathbb{G}/\mathbb{K} is a natural generalization of the q -deformed projective spaces (in particular the standard Podleś sphere $\mathbb{S}_q^2 = \mathbb{CP}^1$). In all cases the unique invariant state under the \mathbb{G} -action is the restriction to $C(\mathbb{S}_{\mathbb{G}})$ of the Haar state on $C(\mathbb{G})$.

6.3 Subproduct systems of \mathbb{G} -representations

The subproduct system associated with a compact quantum group \mathbb{G} will be a subproduct \mathfrak{H}_{\bullet} in which the Hilbert space \mathfrak{H}_m is contained in the m th tensor power $\mathfrak{H}^{\otimes m}$ of the fundamental representation \mathfrak{H} of \mathbb{G} .

The idea is based on the observation that if $u \in M_n(\mathbb{C}) \otimes C(\mathbb{G})$ is a representation of a compact matrix quantum group \mathbb{G} then the first row $z_1 := u_{1,1}, \dots, z_n := u_{1,n}$ of u transforms as the representation u under the “left translation” action λ of \mathbb{G} given by restricting the comultiplication Δ :

$$\lambda(z_k) := \Delta(z_k) = \sum_{j=1}^n z_j \otimes u_{j,k}.$$

The constructions below can easily be made more general but we shall always assume that u is irreducible. In fact we shall, for simplicity and concreteness, from now on assume that u is the defining representation of \mathbb{G} . Let \mathfrak{H} be the n -dimensional Hilbert space with basis vectors z_1, \dots, z_n . In most cases there are subrepresentations of \mathbb{G} contained in the tensor product $\mathfrak{H} \otimes \mathfrak{H}$. Keeping only the largest \mathbb{G} -invariant subspace \mathfrak{H}_2 of $\mathfrak{H} \otimes \mathfrak{H}$ we obtain another irreducible representation $u^{(2)}$ of \mathbb{G} . Indeed, \mathfrak{H}_2 can be identified with the span of $z_j z_k$ for $j, k = 1, \dots, n$, and then $u^{(2)}$ is obtained by the restricting the comultiplication, just as for u . Continuing like this we obtain a family $(\mathfrak{H}_m)_{m \in \mathbb{N}_0}$ of Hilbert spaces satisfying the subproduct condition (2.1).

Definition 6.19. Let \mathbb{G} be a compact matrix quantum group. The subproduct system \mathfrak{H}_{\bullet} just described will be referred to as the **the \mathbb{G} -subproduct system**.

If we dropped the requirement that u generates $C(\mathbb{G})$ then Definition 6.19 makes no use of the fact that $C(\mathbb{G})$ is finitely generated, and hence it works for all compact quantum groups. On the other hand, if we do not require irreducibility but \mathbb{G} is a matrix group, we may always find a *self-conjugate* unitary finite-dimensional representation u whose coefficients generate $C(\mathbb{G})$. Indeed, if u generates $C(\mathbb{G})$ then so does $u \oplus \bar{u}$, and the latter is self-conjugate. If u is self-conjugate and generates $C(\mathbb{G})$, every irreducible representation of \mathbb{G} is obtained as \mathfrak{H}_m for a unique $m \in \mathbb{N}_0$. However, for definiteness we shall always suppose that \mathbb{G} is a compact *matrix* quantum group and that u is the defining representation, assumed irreducible.

Example 6.20. For $\mathbb{G} = G$ a classical compact Lie group, it is well known that the representation $\mathfrak{H}_{\lambda+\mu}$ with dominant weight $\lambda + \mu$ occurs exactly once in the tensor product $\mathfrak{H}_{\lambda} \otimes \mathfrak{H}_{\mu}$; this is the “Cartan product” of \mathfrak{H}_{λ} and \mathfrak{H}_{μ} [East1]. In particular, if \mathfrak{H}_{\bullet} denotes the G -subproduct system then \mathfrak{H}_{m+1} is the Cartan product of \mathfrak{H}_m and \mathfrak{H} . This subproduct system is commutative, i.e. $\mathfrak{H}_{\bullet} \subseteq \mathfrak{H}^{\vee \bullet}$, and the associated projective variety mentioned in §3.1 is a coadjoint orbit, isomorphic to G/K for some closed subgroup K of G .

Example 6.21. The $\mathrm{SU}(n)$ -subproduct system coincides with the fully symmetric subproduct system $\mathfrak{H}^{\vee \bullet}$ (Example 2.2).

Example 6.22. The \mathbb{G} -subproduct system of the universal quantum group $\mathbb{G} = A_u(Q)$ with positive $Q \in \mathrm{GL}(n, \mathbb{C})$ is the product system $\mathfrak{H}^{\otimes \bullet}$ because each power $u^{\otimes m}$ of the defining representation is irreducible [Ban4].

Example 6.23. For $\mathbb{G} = B_u(F)$ we have self-conjugacy $u \simeq \bar{u}$ [Ban3]. So if u is irreducible then, as mentioned above, every $\lambda \in \mathrm{Irrep} \mathbb{G}$ occurs as $\mathfrak{H}_\lambda \cong \mathfrak{H}_m$ for some $m \in \mathbb{N}_0$.

The \mathfrak{H}_m 's are all irreducible and usually pairwise inequivalent. The only examples where they are not all inequivalent are those where $C(\mathbb{G})$ is finite-dimensional.

6.4 Berezin quantization of $C(\mathbb{G}/\mathbb{K})$

6.4.1 Covariant symbols as first-row matrix coefficients

Recall the formula (4.9) for the covariant symbol derived in Proposition 4.19. We now observe that for any compact matrix quantum groups \mathbb{G} , the same formula (with u now being the defining representation of \mathbb{G}) appears when taking matrix coefficients of the representation $\mathrm{Ad}(u^{(m)}) \simeq u^{(m)} \otimes \bar{u}^{(m)}$ on $\mathcal{B}(\mathfrak{H}_m)$.

Definition 6.24. We say that an element of $C(\mathbb{G}/\mathbb{K})$ is **normally ordered** if all z_j 's occur to the left of the z_k^* 's.

Lemma 6.25. *Suppose that every element in $C(\mathbb{G}/\mathbb{K})$ can be written in normally ordered form. Then the C^* -algebra $C(\mathbb{G}/\mathbb{K})$ can be generated by covariant symbols*

$$\varsigma_{\mathbb{G}}^{(m)}(A) := (\mathrm{Tr} \otimes \mathrm{id})((A \otimes \mathbf{1})u^{(m)c*}(|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}| \otimes \mathbf{1})u^{(m)c}) \quad (6.9)$$

for all $A \in \mathcal{B}(\mathfrak{H}_m)$ and all $m \in \mathbb{N}_0$.

Proof. Write $\phi_A := \mathrm{Tr}(A \cdot)$. Then the elements

$$\varsigma_{\mathbb{G}}^{(m)}(A) = (\phi_A \otimes \mathrm{id})(u^{(m)c*}(|e_1^{\otimes m}\rangle\langle e_1^{\otimes m}| \otimes \mathbf{1})u^{(m)c}), \quad A \in \mathcal{B}(\mathfrak{H}_m)$$

form a set which consists exactly of those matrix coefficients of the representation $u^{(m)c*} \otimes u^{(m)c} \simeq u^{(m)} \otimes \bar{u}^{(m)}$ which are contained in $C(\mathbb{G}/\mathbb{K})$. \square

6.4.2 Intertwining the actions

Let W be the multiplicative unitary of \mathbb{G} , i.e. the unitary on $L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$ implementing the comultiplication Δ (cf. §6.1.3). Then the restriction of W to $L^2(\mathbb{G}/\mathbb{K}) \otimes L^2(\mathbb{G})$ identifies with

$$u^{(\mathbb{N})} := (u^{(m)})_{m \in \mathbb{N}_0} \in c_b(\hat{\mathbb{G}}) \otimes C(\mathbb{G}).$$

We have an action of \mathbb{G} on $\mathcal{R}(\mathbb{G})$ given by $\mathrm{Ad}(W)$ (i.e. the same formula as for Δ but now applied to elements of a different algebra; of course this action extends to all of $\mathcal{B}(L^2(\mathbb{G}))$). The restriction of this \mathbb{G} -action to $\mathcal{B}(\mathfrak{H}_{\mathbb{N}})$ is just $\mathrm{Ad}(u^{(\mathbb{N})})$. Similarly, the extension of $\hat{\Delta}$ to $\mathcal{B}(L^2(\mathbb{G}))$ restricts to a right $\hat{\mathbb{G}}$ -action on $L^\infty(\mathbb{G})$.

So we have the following actions of \mathbb{G} and $\hat{\mathbb{G}}$ on $C(\mathbb{G}/\mathbb{K})$.

Definition 6.26. The left \mathbb{G} -action $\alpha^{\mathbb{G}} : C(\mathbb{G}/\mathbb{K}) \rightarrow C(\mathbb{G}/\mathbb{K}) \otimes C(\mathbb{G})$ is defined by

$$\alpha^{\mathbb{G}}(\varsigma_{\mathbb{G}}^{(m)}(A)) := (\varsigma_{\mathbb{G}}^{(m)} \otimes \mathrm{id})(u^{(m)}(A \otimes \mathbf{1})u^{(m)*})$$

for all $A \in \mathcal{B}(\mathfrak{H}_m)$ and all $m \in \mathbb{N}_0$. The right $\hat{\mathbb{G}}$ -action $\hat{\alpha}^{\mathbb{G}} : C(\mathbb{G}/\mathbb{K}) \rightarrow c_b(\hat{\mathbb{G}}) \otimes C(\mathbb{G}/\mathbb{K})$ is defined by

$$\hat{\alpha}^{\mathbb{G}}(\varsigma_{\mathbb{G}}^{(m)}(A)) := (\mathrm{id} \otimes \varsigma_{\mathbb{G}}^{(m)})(u^{(m)*}(\mathbf{1} \otimes A)u^{(m)}).$$

Lemma 6.27. *The map $\varsigma_{\mathbb{G}}^{(m)}$ is \mathbb{G} - and $\hat{\mathbb{G}}$ -equivariant,*

$$\Delta \circ \varsigma_{\mathbb{G}}^{(m)}(A) = (\varsigma_{\mathbb{G}}^{(m)} \otimes \text{id})(u^{(m)}(A \otimes \mathbf{1})u^{(m)*}), \quad (6.10)$$

$$\hat{\Delta} \circ \varsigma_{\mathbb{G}}^{(m)}(A) = (\text{id} \otimes \varsigma_{\mathbb{G}}^{(m)})(u^{(m)*}(\mathbf{1} \otimes A)u^{(m)}). \quad (6.11)$$

The actions $\alpha^{\mathbb{G}}$ and $\hat{\alpha}^{\mathbb{G}}$ therefore coincide on $C(\mathbb{G}/\mathbb{K}) \subset L^\infty(\mathbb{G})$ with the left regular action of \mathbb{G} and the right regular action of $\hat{\mathbb{G}}$, respectively.

Proof. Let $\alpha^{(m)}(A) := u^{(m)}(A \otimes \mathbf{1})u^{(m)*}$ and $\hat{\alpha}^{(m)}(A) := u^{(m)*}(\mathbf{1} \otimes A)u^{(m)}$. We use the defining property of a left action,

$$(\alpha^{(m)} \otimes \text{id}) \circ \alpha^{(m)} = (\text{id} \otimes \Delta) \circ \alpha^{(m)}.$$

Formula (4.9) gives

$$\begin{aligned} (\varsigma_{\mathbb{G}}^{(m)} \otimes \text{id})(\alpha^{(m)}(A)) &= (\text{Tr} \otimes \text{id} \otimes \text{id})((\alpha^{(m)} \otimes \text{id}) \circ \alpha^{(m)}(A)(|e_1^m\rangle\langle e_1^m| \otimes \mathbf{1})) \\ &= (\text{Tr} \otimes \text{id} \otimes \text{id})((\text{id} \otimes \Delta) \circ \alpha^{(m)}(A)(|e_1^m\rangle\langle e_1^m| \otimes \mathbf{1})) \\ &= \Delta(\text{Tr} \otimes \text{id})(\alpha^{(m)}(A)(|e_1^m\rangle\langle e_1^m| \otimes \mathbf{1})) = \Delta \circ \varsigma_{\mathbb{G}}^{(m)}(A). \end{aligned}$$

The proof of (6.11) is identical, using $(\text{id} \otimes \hat{\alpha}^{(m)}) \circ \hat{\alpha}^{(m)} = (\hat{\Delta} \otimes \text{id}) \circ \hat{\alpha}^{(m)}$. \square

6.4.3 $C(\mathbb{G}/\mathbb{K})$ as an inductive limit

Now we will, for certain compact matrix quantum groups \mathbb{G} , realize the first-row algebra $C(\mathbb{G}/\mathbb{K})$ as a projective limit \mathcal{B}^∞ . From our previous results we have then obtained $C(\mathbb{G}/\mathbb{K})$ as the $U(1)$ -invariant part of the Cuntz–Pimsner algebra of the \mathbb{G} -subproduct system.

Let $\mathbb{G} \subset A_u(Q)$ be a compact matrix quantum group (with faithful Haar measure), with $Q \in GL(n, \mathbb{C})$, and let $C(\mathbb{G}/\mathbb{K})$ be the $U(1)$ -invariant part of the first-row algebra $C(\mathbb{S}_{\mathbb{G}})$.

Let $\mathfrak{H}_\bullet = \mathfrak{H}_\bullet^{\mathbb{G}}$ be the \mathbb{G} -subproduct system and let $\mathcal{O}_{\mathfrak{H}}$ be its Cuntz–Pimsner algebra. In §5 we defined the Toeplitz quantization $\zeta^{(m)}$ as a map from $\mathcal{O}_{\mathfrak{H}}$ to \mathcal{B}^∞ . In that way we could realize an isomorphism between $\mathcal{O}_{\mathfrak{H}}^{(0)} \cong \mathcal{B}_\infty$ and the projective limit \mathcal{B}^∞ . In this section we shall use the same strategy but with a map

$$\zeta_{\mathbb{G}}^{(m)} : C(\mathbb{G}/\mathbb{K}) \rightarrow \mathcal{B}(\mathfrak{H}_m), \quad (6.12)$$

i.e. we quantize $C(\mathbb{G}/\mathbb{K})$ instead of $\mathcal{O}_{\mathfrak{H}}^{(0)}$. We define (6.12) to be the adjoint of the map $\varsigma_{\mathbb{G}}^{(m)}$ appearing in (6.9). We shall in this way obtain an isomorphism between $C(\mathbb{G}/\mathbb{K})$ and \mathcal{B}^∞ , and hence, due to $\mathcal{B}_\infty \cong \mathcal{B}^\infty$, we will arrive at the following result.

Theorem 6.28. *Assume that normal ordering is possible in $C(\mathbb{G}/\mathbb{K})$. Then there is a C^* -isomorphism between $C(\mathbb{G}/\mathbb{K})$ and the $U(1)$ -invariant part of the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ of the \mathbb{G} -subproduct system:*

$$C(\mathbb{G}/\mathbb{K}) \cong \mathcal{O}_{\mathfrak{H}}^{(0)}.$$

This isomorphism is given by the total Toeplitz map $\zeta_{\mathbb{G}} = \prod_m \zeta_{\mathbb{G}}^{(m)}$, and it intertwines the ergodic \mathbb{G} - and $\hat{\mathbb{G}}$ -actions as well as the \mathbb{G} -invariant states.

Remark 6.29 (Normal ordering). For general \mathbb{G} , the map $\zeta_{\mathbb{G}}$ maps the normally ordered part of $C(\mathbb{G}/\mathbb{K})$ onto $\mathcal{O}_{\mathfrak{H}}^{(0)}$. For instance, normal ordering is possible for $\mathbb{G} = B_u(F)$ but not for $A_u(Q)$. In fact, $A_u(Q)$ is our only example where $\zeta_{\mathbb{G}}$ is not an isomorphism. Commutative fusion rules implies normal ordering (recall Definition 6.5).

So let us begin by defining \mathcal{B}^∞ to be the projective limit of the system $(\mathcal{B}(\mathfrak{H}_\bullet), j_{\bullet, \bullet})$, where \mathfrak{H}_\bullet is the \mathbb{G} -subproduct system. Here the operator Q on \mathfrak{H} which appears in the construction of \mathcal{B}^∞ (§4.5) is taken to be the same as the matrix defining $A_u(Q) \supset \mathbb{G}$, assuming Q is equal to its transpose. Thus, $\mathcal{B}(\mathfrak{H}_m)$ is equipped with the ϕ_m -inner product, where $\phi_m = \text{Tr}(\rho_Q^{(m)} \cdot)$ is the state defined by

the density matrix $\rho_Q^{(m)} := Q_m / \text{Tr}(Q_m)$. We may therefore regard \mathcal{B}^∞ as a subset of $c_b(\hat{\mathbb{G}})$ having trivial intersection with $c_0(\hat{\mathbb{G}})$.

From our general results we have an isomorphism $\zeta(\mathcal{B}_\infty) \cong \mathcal{B}_\infty$ which realizes the projective limit as an inductive limit. We stress again that this isomorphism ζ is not the same as the map $\zeta_{\mathbb{G}}$ which we now try to prove is an isomorphism.

The matrix coefficients of the operator $\zeta^{(m)}(f)$ are of the form

$$h(z_{\mathbf{j}} z_{\mathbf{k}}^* f), \quad \mathbf{j}, \mathbf{k} \in \mathbb{F}_n^+ \text{ with } |\mathbf{j}| = m = |\mathbf{k}|. \quad (6.13)$$

Proof of Theorem 6.28. Since the “coefficient map” $\zeta_{\mathbb{G}}^{(m)}$ in (6.9) is injective, its adjoint $\zeta_{\mathbb{G}}^{(m)}$ is surjective. As in the case of \mathcal{B}_∞ , we get that the image of $L^\infty(\mathbb{G}/\mathbb{K})$ under $\zeta_{\mathbb{G}}$ is exactly \mathcal{B}^∞ as a set.

We cannot use the reasoning in the proof of Lemma 5.14 to deduce that $\zeta_{\mathbb{G}}$ is an injection of $C(\mathbb{G}/\mathbb{K})$ into the operator system \mathcal{B}^∞ . On the other hand, we see directly that if $\zeta(f) = 0$ then, using that the matrix coefficients of $\zeta(f)$ are given by (6.13) for all $m \in \mathbb{N}$, we get that f must be orthogonal to the whole normally ordered part of $C(\mathbb{G}/\mathbb{K}) \subset L^2(\mathbb{G})$. Since we have assumed that each $f \in C(\mathbb{G}/\mathbb{K})$ can be normally ordered and that the Haar state is faithful, this means that $f = 0$.

Moreover, $\pi^{-1}(\zeta(\mathcal{B}_\infty))$ is again equal to $\mathcal{T}_{\mathfrak{H}}^{(0)}$. Namely, the proof in §5.7 carries over completely.

As before we get that $\zeta^{(m)} = j_{\infty, m}$. Since we know that $\zeta(\mathcal{B}_\infty)$ is a C^* -algebra (using that it is a quotient of the Toeplitz algebra) with a unique multiplication, we obtain the von Neumann condition, i.e. $\zeta^{(m)}$ is asymptotically a homomorphism (see Corollary 5.24). Thus ζ is an isomorphism for the projective-limit multiplication.

We also know that $\zeta_{\mathbb{G}}$ intertwines the vacuum state $\hat{\varepsilon}$ with the Haar state h restricted to $C(\mathbb{G}/\mathbb{K})$. Composing with the isomorphism $\zeta : \mathcal{B}_\infty \rightarrow \zeta(\mathcal{B}_\infty)$ we get that h is intertwined with the limit state ω_Q . Finally, Lemma 6.27 shows that $\zeta_{\mathbb{G}}$ is \mathbb{G} - $\hat{\mathbb{G}}$ -equivariant. \square

6.4.4 $C(\mathbb{S}_{\mathbb{G}})$ as an inductive limit

Corollary 6.30. *Let \mathbb{G} be a compact matrix quantum group with faithful Haar measure $h : C(\mathbb{G}) \rightarrow \mathbb{C}$ such that normal ordering is possible in $C(\mathbb{G}/\mathbb{K})$. Then there is a \mathbb{G} - $\hat{\mathbb{G}}$ -equivariant isomorphism between the first-row algebra $C(\mathbb{S}_{\mathbb{G}})$ and the Cuntz–Pimsner algebra $\mathcal{O}_{\mathfrak{H}}$ of the \mathbb{G} -subproduct system,*

$$C(\mathbb{S}_{\mathbb{G}}) \cong \mathcal{O}_{\mathfrak{H}}.$$

In particular, $\mathcal{O}_{\mathfrak{H}}$ carries an ergodic action of \mathbb{G} in which each irreducible representation of \mathbb{G} occurs exactly once.

Proof. For notation simplicity we identify $\mathcal{O}_{\mathfrak{H}}^{(0)}$ with the inductive limit \mathcal{B}_∞ and the modules $\mathcal{O}_{\mathfrak{H}}^{(k)}$ with the modules $\mathcal{E}^{(k)}$.

Since $\mathcal{B}_\infty \cong C(\mathbb{G}/\mathbb{K})$, we know that there is a basis e_1, \dots, e_n for \mathfrak{H} such that the Q -sphere condition (6.4) is satisfied by the generators Z_1, \dots, Z_n of $\mathcal{O}_{\mathfrak{H}}$, just as it is for the generators z_1, \dots, z_n of $C(\mathbb{S}_{\mathbb{G}})$. This says precisely that Z_1, \dots, Z_n and $Q_{1,1}^{-1/2} Z_1, \dots, Q_{n,n}^{-1/2} Z_n$ are standard right and left tight normalized frames for the \mathcal{B}_∞ -bimodule $\mathcal{E}^{(1)}$, respectively; for all $\xi \in \mathcal{E}^{(1)}$,

$$\sum_{k=1}^n \langle \xi | Z_k \rangle_{\text{right}} \langle Z_k | \xi \rangle_{\text{right}} = \langle \xi | \xi \rangle_{\text{right}},$$

$$\sum_{k=1}^n \langle \xi | Q_{k,k}^{-1/2} Z_k \rangle_{\text{left}} \langle Q_{k,k}^{-1/2} Z_k | \xi \rangle_{\text{left}} = \sum_{k=1}^n \xi Q_{k,k}^{-1} Z_k^* Z_k \xi^* = \xi \xi^* = \langle \xi | \xi \rangle_{\text{left}},$$

and identically for $C(\mathbb{S}_{\mathbb{G}})^{(1)}$ and the z_j ’s. If we identify $C(\mathbb{G}/\mathbb{K})$ with \mathcal{B}_∞ , this means that the projection $P^{(1)} \in M_n(\mathbb{C}) \otimes C(\mathbb{G}/\mathbb{K})$ which defines the module $C(\mathbb{S}_{\mathbb{G}})^{(1)}$ coincides with the projection

which defines the module $\mathcal{E}^{(1)}$. So the modules are the same and the isomorphism $C(\mathbb{S}_{\mathbb{G}}) \cong \mathcal{O}_{\mathfrak{H}}$ is clear.

For the \mathbb{G} - $\hat{\mathbb{G}}$ -equivariance, we must first define actions on $\mathcal{O}_{\mathfrak{H}}$. But since we know that $C(\mathbb{S}_{\mathbb{G}}) \cong \mathcal{O}_{\mathfrak{H}}$ we can just specify these action on generators Z_1, \dots, Z_n by the same formulas as for $C(\mathbb{S}_{\mathbb{G}})$.

The last statement is due to Lemma 6.18. \square

6.5 Comparison with Poisson and Martin boundaries

6.5.1 Poisson integral versus total Toeplitz map

Let \mathbb{G} be a compact matrix quantum group with commutative fusion rules (see Definition 6.5) and faithful Haar measure. The Poisson boundary to be discussed here is the one defined in [Iz1], so if we were phrasing things in terms of random walks (we shall not), there would in the background be a representation u of \mathbb{G} whose coefficients generate $C(\mathbb{G})$ (without any need of the adjoints $u_{j,k}^*$).

Izumi defines [Iz1, Lemma 3.8] the **Poisson integral** to be the unital completely positive map $\Theta : L^\infty(\mathbb{G}) \rightarrow \mathcal{R}(\mathbb{G})$ given by

$$\Theta(f) := (\text{id} \otimes h)(W^*(\mathbf{1} \otimes f)W), \quad (6.14)$$

where W is the fundamental unitary (§6.1.3). Similar to the projective limit \mathcal{B}^∞ which is the image of our total Toeplitz map ζ , the image of map Θ is an operator system, usually denoted by $H^\infty(\hat{\mathbb{G}})$, which can be made into a von Neumann algebra by replacing the operator multiplication by the new one. Moreover, Θ is a complete order isomorphism onto its image. A possible definition of the **Poisson boundary** of $\hat{\mathbb{G}}$ is then as the preimage, say $L^\infty(\mathbb{G}/\mathbb{T})$, of Θ in $L^\infty(\mathbb{G})$. We then refer to the abstract object \mathbb{G}/\mathbb{T} as the Poisson boundary of $\hat{\mathbb{G}}$. The notation \mathbb{G}/\mathbb{T} is chosen to indicate that \mathbb{G} and $\hat{\mathbb{G}}$ act ergodically on $L^\infty(\mathbb{G}/\mathbb{T})$.

The Poisson boundary \mathbb{G}/\mathbb{T} is defined in terms of a von Neumann algebra. In order to compare \mathbb{G}/\mathbb{T} with what we have denoted \mathbb{G}/\mathbb{K} , note that ζ extends to a normal completely positive map (denoted by the same symbol)

$$\zeta : L^\infty(\mathbb{G}/\mathbb{K}) \rightarrow \mathcal{B}^\infty \quad (6.15)$$

and this is the “first-row” version of the Poisson integral. Using it one can carry out Berezin quantization on the level of von Neumann algebras. Inspiring work here is [INT1].

The Poisson integral (6.14) can be decomposed into components $\Theta_\lambda : L^\infty(\mathbb{G}) \rightarrow \mathcal{B}(\mathfrak{H}_\lambda)$ for $\lambda \in \text{Irrep } \mathbb{G}$, and doing so one easily calculates the adjoints $\Theta_\lambda^* : \mathcal{B}(\mathfrak{H}_\lambda) \rightarrow L^\infty(\mathbb{G})$. Noticing the similarity to Berezin quantization, [INT1] referred to the composition $\Theta_\lambda^* \circ \Theta_\lambda$ as the “Berezin transform”. This terminology is not entirely fortunate because $\Theta_\lambda^* \circ \Theta_\lambda$ does not coincide with the usual notion of Berezin transform when $\mathbb{G} = G$ is an ordinary group. The issue is that Θ_λ^* is obtained by tracing against the invertible operator Q_λ (which is of full rank) instead of a rank-1 projection. The distinction is the use of “first-row” versus all of \mathbb{G} . This distinction persists even if we, as Izumi does, assume that every irreducible representation of \mathbb{G} is contained in some power of u .

It is therefore interesting that the final results (\mathbb{G}/\mathbb{K} and \mathbb{G}/\mathbb{T}) are not very different. For $\text{SU}_q(2)$ they even coincide. In general, we should view \mathbb{G}/\mathbb{K} as a (noncommutative) non-maximal flag variety (prototype example being \mathbb{CP}_q^{n-1}) while \mathbb{G}/\mathbb{T} is the maximal flag variety (so \mathbb{T} is the “maximal torus”); cf. [Tom1].

The transition between classical and quantum Poisson boundaries is rather involved [NT1]. In fact, if $\mathbb{G} = G$ is an ordinary compact group then the Poisson boundary is trivial: $L^\infty(\mathbb{G}/\mathbb{T}) = \mathbb{C}\mathbf{1}$ [Iz1, Cor. 3.9]. In contrast, Berezin quantization carries over in perfect analogy with the commutative case.

6.5.2 Markov operator

The set $H^\infty(\Phi)$ of fixed points of a normal completely positive map Φ on a von Neumann algebra is an ultraweakly closed operator system which can be made into a von Neumann algebra by replacing the operator multiplication by the so-called “Choi–Effros multiplication” [Ar10, Thm. 3.1], [Iz4].

The new multiplication on the Poisson boundary $H^\infty(\hat{\mathbb{G}})$ mentioned above is just an example of a Choi–Effros multiplication. The completely positive map on $\mathcal{R}(\mathbb{G})$ whose fixed-point set equals $H^\infty(\hat{\mathbb{G}})$ takes the role of Markov operator for the “noncommutative random walk” on $\hat{\mathbb{G}}$.

The following can be summarized by saying that with Berezin quantization one ends up with a random walk on the “dual” of \mathbb{G}/\mathbb{K} instead of the dual of \mathbb{G} . Note however that it works for any subproduct system \mathfrak{H}_\bullet . Fix thus a subproduct system \mathfrak{H}_\bullet and denote as usual by $\Gamma_b = \Gamma_b(\mathcal{B}_\bullet)$ the von Neumann-algebraic direct sum of the matrix algebras $\mathcal{B}_m := \mathcal{B}(\mathfrak{H}_m)$.

Definition 6.31. The **Markov operator** on \mathcal{B}_\bullet is the unital normal completely positive map $\Phi : \Gamma_b \rightarrow \Gamma_b$ defined by

$$\Phi(X_\bullet) := X_{\bullet-1} = (j_{m,m-1}(X_m))_{m \in \mathbb{N}}.$$

Proposition 6.32. *The set $H^\infty(\Phi)$ of Φ -fixed points in $\Gamma_b(\mathcal{B}_\bullet)$ is equal to the projective limit \mathcal{B}^∞ . In particular, \mathcal{B}^∞ is a von Neumann algebra. On the subset $\zeta(\mathcal{B}_\infty) \subset \mathcal{B}^\infty$, the Choi–Effros multiplication coincides with the projective-limit multiplication.*

Proof. The first statement is clear, so \mathcal{B}^∞ is a von Neumann algebra. To prove the last statement we use the result [Iz3, Cor. 5.2] that the Choi–Effros product of $X, Y \in H^\infty(\Phi)$ is given by

$$X \diamond Y = \lim_{r \rightarrow \infty} \Phi^r(XY),$$

where XY is the multiplication in $\mathcal{B}(\mathfrak{H}_\mathbb{N})$ and the limit is in the strong operator topology. Now, the m th component of $X \diamond Y$ is

$$\begin{aligned} (X \diamond Y)_m &= \left(\lim_{r \rightarrow \infty} \Phi^r(XY) \right)_m = \lim_{r \rightarrow \infty} j_{m+r,m}((XY)_{m+r}) \\ &= j_{\infty,m} \left(\lim_{r \rightarrow \infty} (XY)_{m+r} \right) \\ &= \left(\lim_{l \rightarrow \infty} \zeta^{(l)}(XY) \right)_m, \end{aligned}$$

where we used that $X_m = j_{\infty,m}(X) = \zeta^{(m)}(X)$. This shows that \diamond is the projective-limit multiplication (5.15) whenever we have convergence in norm. Since norm-convergence holds for $X = \zeta(f)$ and $Y = \zeta(g)$ with $f, g \in \mathcal{B}_\infty$, the claim holds. \square

6.5.3 Martin boundaries

While the Poisson boundary is a measure-theoretic object defined via a von Neumann algebra, the Martin boundary is specified in terms of a C^* -algebra [NT1]. Its relation to Berezin quantization is the same “first-row versus all-of- \mathbb{G} ” story as with the Poisson boundary but we shall discuss only a special case in which \mathbb{G}/\mathbb{K} agrees with the Martin boundary of $\hat{\mathbb{G}}$. The reason for this coincidence is that the defining representation of the chosen \mathbb{G} is self-conjugate and irreducible, so that \mathfrak{H}_\bullet contains all irreducible representations.

Our approach here via inductive limits was partially inspired by [VVer1], where they construct a “Martin boundary” of the dual of \mathbb{G} for $\mathbb{G} = B_u(F)$ in the same way. Our notation \mathcal{B}_∞ is chosen to make comparison with that paper easy. Let $F \in \text{GL}(n, \mathbb{C})$ such that $\bar{F}F = \pm \mathbf{1}$; this ensures that the defining representation of $\mathbb{G} = B_u(F)$ is irreducible. By construction, the Martin boundary of $\hat{\mathbb{G}}$ is equal to the inductive limit \mathcal{B}_∞ of the \mathbb{G} -subproduct system.

In [VaVe1], another realization of the Martin boundary was accomplished. First define $\mathcal{B}_u(F, F_q)$ to be the universal C^* -algebra generated by the entries of a unitary $2 \times n$ matrix Y satisfying

$$Y = F_q Y^c F^{-1}. \quad (6.16)$$

where $F_q := \begin{pmatrix} 0 & |q|^{1/2} \\ \pm |q|^{-1/2} & 0 \end{pmatrix}$, with q defined by $|q + q^{-1}| = \text{Tr}(F^*F)$ and $F\bar{F} = \pm q$ and we wrote $q = \mp |q|$. It is shown in [VaVe1] that the $\text{U}(1)$ -action ρ on $\mathcal{B}_u(F, F_q)$ given by

$$\rho_\lambda(Y) := \begin{pmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{pmatrix} Y, \quad \forall \lambda \in \mathbb{C}, |\lambda| = 1$$

allows recovering the Martin boundary as the fixed-point algebra $\mathcal{B}_u(F, F_q)^{U(1)}$.

Since our inductive limit \mathcal{B}_∞ coincides with the Martin boundary of the dual of $B_u(F)$, we know (using Theorem 6.28) that $\mathcal{B}_u(F, F_q)^{U(1)}$ must coincide with $C(\mathbb{G}/\mathbb{K})$. This can be seen directly. Let z_1, \dots, z_n and w_1, \dots, w_n be the elements of the first and second row of Y respectively. Then (6.16) reads

$$z_k = |q|^{1/2} \sum_{s=1}^n w_s^* F_{s,k}^{-1}, \quad \forall k = 1, \dots, n. \quad (6.17)$$

The action ρ_z is given by

$$\rho_\lambda(z_k) = \lambda z_k, \quad \rho_\lambda(w_k) = \bar{\lambda} w_k,$$

so the fixed-point algebra consists of elements of the form $z_j w_k$, and their adjoints, as well as $z_j z_k^*$ and $w_j w_k^*$. Now (6.17) shows that $\mathcal{B}_u(F, F_q)^{U(1)} = C(\mathbb{G}/\mathbb{K})$, as asserted.

Note however that $\mathcal{B}_u(F, F_q)$ is not at all the same as the first-row algebra $C(\mathbb{S}_\mathbb{G})$.

Thus we have one example where the Martin boundary of a discrete quantum group identifies with the object \mathbb{G}/\mathbb{K} defined in this paper. Also, let $q \in (0, 1]$. Then for the $\mathbb{G} = \mathrm{SU}_q(2)$ -subproduct system \mathfrak{H}_\bullet we have an equivariant isomorphism

$$\mathcal{O}_{\mathfrak{H}}^{(0)} \cong C(\mathbb{S}_q^2),$$

where $\mathbb{G}/\mathbb{K} = \mathbb{S}_q^2$ is the Podleś sphere. For $q < 1$, this shows again that our inductive limit \mathcal{B}_∞ coincides with the C^* -algebra referred to as the Martin boundary of the dual of $\mathrm{SU}_q(2)$ in [VVer1] (because the same boundary is known to be the Podleś sphere [NT1]). For $q = 1$, we have agreement with Biane’s Martin boundary of the dual of $\mathrm{SU}(2)$ [Biane1].

7 Concluding remarks

We have seen that the inductive limit \mathcal{B}_∞ associated with a subproduct system \mathfrak{H}_\bullet is a sensible generalization of the C^* -algebra $C(M)$ of continuous functions on a quantizable Kähler manifold, in the case the Kähler structure is projectively induced (so M is embeddable as a submanifold of projective space). We may therefore write

$$C(\mathbb{M}) := \mathcal{B}_\infty,$$

and say that \mathbb{M} is the **noncommutative projective variety** associated to \mathfrak{H}_\bullet . We may also refer to elements of

$$C(\mathbb{S}_\mathbb{M}) := \mathcal{O}_{\mathfrak{H}}.$$

as functions on the total space $\mathbb{S}_\mathbb{M}$ of a noncommutative circle bundle over \mathbb{M} . Thus the notation $\mathbb{M} := \mathbb{G}/\mathbb{K}$ and $\mathbb{S}_\mathbb{M} := \mathbb{S}_\mathbb{G}$ would be consistent with that in §6 when \mathfrak{H}_\bullet is the \mathbb{G} -subproduct system. In [An5] we refer to \mathbb{M} as the “dequantization manifold”.

By defining $C(\mathbb{M})$ to be *equal* (and not just isomorphic) to the inductive limit \mathcal{B}_∞ , the noncommutative space \mathbb{M} comes with more structure than just its topology. Namely, if $\mathbb{M} = M$ is commutative then the inductive system gives an embedding into projective space \mathbb{CP}^{n-1} and, if M is non-singular, endows M with a complex-analytic (in particular smooth) structure, a polarization L (choice of ample line bundle), an inner product on $H^0(M; L)$ (the one we started with, making $H^0(M; L)$ into the Hilbert space \mathfrak{H}), a Hermitian metric on L (the one defining the $*$ -structure on $\mathcal{O}_\mathfrak{H}$; this is just the Fubini–Study metric associated with the inner product on \mathfrak{H}) and a volume form on M (viz. the limit state, which need not be the same as Fubini–Study volume form). As we have seen, these structures have perfect generalizations to the noncommutative setting. Note that the quantum homogeneous spaces \mathbb{G}/\mathbb{K} are “balanced” in the sense that the limit state on $C(\mathbb{G}/\mathbb{K})$ coincides with the state induced by the Haar state on $C(\mathbb{G})$.

The covariant symbols $\zeta^{(m)}(A)$ and the Toeplitz operators $\zeta^{(m)}(f)$ can be expressed in terms of the projections $P^{(m)} \in \mathcal{B}(\mathfrak{H}_m) \otimes C(\mathbb{M})$ which define the modules $\mathcal{E}^{(m)}$. In this way one generalizes

the Rawsnelly coherent-state projections in [RCG1], and in particular the coherent states in [Per1]. We will use this when we discuss the (fuzzy) geometry of \mathbb{M} in another paper.

There are also projections $P^{(-m)}$ and maps $\varsigma^{(-m)}, \xi^{(-m)}$ etc. associated with the modules $\mathcal{E}^{(-m)}$. In this way one quantizes instead the anti-normal part of $C(\mathbb{G}/\mathbb{K})$.

8 References

- [An4] Andersson A. Detailed balance as a quantum-group symmetry of Kraus operators. arXiv:1506.00411 (2015).
- [An5] Andersson A. Dequantization via quantum channels. Lett. Math. Phys. Vol 106, Issue 10, pp. 1397-1414 (2016).
- [AraM1] Ara P, Mathieu M. Local multipliers of C^* -algebras. Springer (2003).
- [Ar6] Arveson W. Subalgebras of C^* -algebras III: Multivariable operator theory. Acta Math. Vol 181, pp. 159-228 (1998).
- [Ar8] Arveson W. p -Summable commutators in dimension d . J. Operator Theory. Vol 54, pp. 101-117 (2005).
- [Ar10] Arveson W. Noncommutative Poisson boundaries. Unpublished. Available at <http://math.berkeley.edu/~arveson/Dvi/290F04/22Sept.pdf> (2004).
- [BDLMC] Balachandran AP, Dolan BP, Lee JH, Martin X, O'Connor D. Fuzzy complex projective spaces and their star-products. J. Geom. Phys. Vol 43, pp. 184 (2002).
- [Ban3] Banica T. Théorie des représentations du groupe quantique compact libre $O(n)$. C. R. Acad. Sci. Paris Sér. I Math. Vol 322, pp. 241-244 (1996).
- [Ban4] Banica T. Le groupe quantique compact libre $U(n)$. Comm. Math. Phys. Vol 190, pp. 143-172 (1997).
- [BaGo1] Banica T, Goswami D. Quantum isometries and noncommutative spheres. Comm. Math. Phys. Vol 298, Issue 2, pp. 343-356 (2010).
- [BMT1] Bédos E, Murphy GJ, Tuset L. Co-amenability of compact quantum groups. J. Geom. Phys. Vol 40, Issue 2, pp. 129-153 (2001).
- [BeSl1] Berceanu S, Schlichenmaier M. Coherent state embeddings, polar divisors and Cauchy formulas. J. Geom. Phys. Vol, Issue 34, pp. 336-358 (2000).
- [Bere1] Berezin FA. Quantization. Math. USSR-Izv. Vol 38, pp. 1116-1175 (1974).
- [Bere2] Berezin FA. General concept of quantization. Comm. Math. Phys. Vol 40, pp. 153-174 (1995).
- [Bere3] Berezin FA. Covariant and contravariant symbols of operators. Math. USSR-Izv. Vol 6, Issue 5, p. 1117 (1972).
- [Bere4] Berezin FA. Some remarks about the associative envelope of a Lie algebra. Funct. Anal. Appl. Vol 1, pp. 91-102 (1967).
- [Biane1] Biane P. Introduction to random walks on noncommutative spaces. In: Quantum Potential Theory. pp. 61-116. Springer (2008).
- [Bied1] Biedenharn LC. The quantum group $SU_q(2)$ and a q -analogue of the boson operators. J. Phys. A: Math. Gen. Vol 22, pp. L873-L878 (1989).
- [Blac1] Blackadar B. Operator algebras: theory of C^* -algebras and von Neumann algebras. Springer (2006).
- [BKil1] Blackadar B, Kirchberg E. Generalized inductive limits of finite-dimensional C^* -algebras. Math. Ann. Vol 307, pp. 343-380 (1997).
- [BMS] Bordemann M, Meinrenken E, Schlichenmaier M. Toeplitz quantization of Kähler manifolds and $\mathfrak{gl}(n), n \rightarrow \infty$ limits. Comm. Math. Phys. Vol 165, Issue 2, pp. 281-296 (1994).
- [BHSS] Bordemann M, Hoppe J, Schaller P, Schlichenmaier M. $\mathfrak{gl}(\infty)$ and geometric quantization. Comm. Math. Phys. Vol 138, Issue 2, pp. 209-244 (1991).

- [BLY1] Bourguignon J-P, Li P, Yau ST. Upper bound for the first eigenvalue of algebraic submanifolds. *Comment. Math. Helv.* Vol 69, pp. 199-207 (1994).
- [BrOz1] Brown NP, Ozawa N. C^* -algebras and finite-dimensional approximations. *Amer. Math. Soc.* (2008).
- [CGR1] Cahen M, Gutt S, Rawnsley J. Quantization of Kähler manifolds I: geometric interpretation of Berezin’s quantization. *J. Geom. Phys.* Vol 7, Issue 1 (1990).
- [CGR2] Cahen M, Gutt S, Rawnsley J. Quantization of Kähler manifolds II. *Trans. Amer. Math.* Vol 337, Issue 1 (1993).
- [CaKe1] Cao HD, Keller J. On the Calabi problem: a finite-dimensional approach. *J. Eur. Math. Soc. (JEMS)* Vol 15, Issue 3, pp. 1033-1065 (2013).
- [DDL1] D’Andrea F, Dabrowski L, Landi G. The noncommutative geometry of the quantum projective plane. *Reviews in Mathematical Physics.* Vol 20, Issue 08, pp. 979-1006 (2008).
- [DRS1] Davidson KR, Ramsey C, Shalit OM. The isomorphism problem for some universal operator algebras. *Adv. Math.* Vol 228, Issue 1, pp. 167-218 (2011).
- [DRS2] Davidson KR, Ramsey C, Shalit OM. Operator algebras for analytic varieties. *Trans. Amer. Math. Soc.* Vol 367, Issue 2, pp. 1121-1150 (2015).
- [Don1] Donaldson SK. Scalar curvature and projective embeddings, I *J. Differential Geom.* Vol 59, pp. 479-522 (2001).
- [Don2] Donaldson SK. Scalar curvature and projective embeddings, II. *Quart. J. Math.* Vol 56, pp. 345-356 (2005).
- [Don3] Donaldson SK. Some numerical results in complex differential geometry. *Pure Appl. Math. Q.* Vol 5, Issue 2, pp. 571-618 (2009).
- [DoWa2] Douglas RG, Wang K. Geometric Arveson–Douglas conjecture and holomorphic extension. *arXiv:1511.00782v1* (2015).
- [East1] Eastwood M. The Cartan product. *Bull. Belg. Math. Soc. Simon Stevin.* Vol. 11, Issue 5, pp. 641-651 (2005).
- [EnEs1] Engliš M, Eschmeier J. Geometric Arveson–Douglas conjecture. *Adv. Math.* Vol 274, pp. 606-630 (2015).
- [Ev1] Evans DE. On \mathcal{O}_n . *Publ. RIMS, Kyoto Univ.* Vol 16, pp. 915-927 (1980).
- [Frank1] Frank M. Isomorphisms of Hilbert C^* -modules and $*$ -isomorphisms of related operator C^* -algebras. *Math. Scand.* Vol 80, pp. 313-319 (1997).
- [Haw1] Hawkins E. Quantization of equivariant vector bundles. *Comm. Math. Phys.* Vol 202, Issue 3, pp. 517-546 (1999).
- [Haw2] Hawkins E. Geometric quantization of vector bundles and the correspondence with deformation quantization. *Comm. Math. Phys.* Vol 215, Issue 2, pp. 409-432 (2000).
- [Haw3] Hawkins E. An obstruction to quantization of the sphere. *Comm. Math. Phys.* Vol 283, Issue 3, pp. 675-699 (2008).
- [Huy] Huybrechts D. *Complex geometry. An introduction.* Springer (2005).
- [Iz1] Izumi M. Non-commutative Poisson boundaries and compact quantum group actions. *Adv. Math.* Vol 169, Issue 1, pp. 1-57 (2002).
- [Iz2] Izumi M. Non-commutative Markov operators arising from subfactors. *Operator algebras and applications.* *Adv. Stud. Pure Math.* Vol 38, pp. 201-217 (2004).
- [Iz3] Izumi M. E_0 -semigroups: around and beyond Arveson’s work. *J. Operator Theory.* Vol 68, Issue 2, pp. 335-363 (2012).
- [Iz4] Izumi M. Non-commutative Poisson boundaries, Discrete geometric analysis. *Contemp. Math.* Vol 347, pp. 69-81 (2004).
- [INT1] Izumi M, Neshveyev S, Tuset L. Poisson boundary of the dual of $SU_q(n)$. *Comm. Math. Phys.* Vol 262, Issue 2, pp. 505-531 (2006).

- [KaSh1] Kakariadis E, Shalit O. On operator algebras associated with monomial ideals in noncommuting variables. arXiv:1501.06495 (2015).
- [KeSh1] Kennedy M, Shalit O. Essential normality, essential norms and hyperrigidity. New York J. Math. Vol 18, pp. 877-890 (2012).
- [KlS] Klimyk AU, Schmüdgen K. Quantum groups and their representations. Vol. 552, Springer, Berlin (1997).
- [Lan1] Landsman NP. Mathematical topics between classical and quantum mechanics. Springer (1998).
- [Lan2] Landsman NP. Strict quantization of coadjoint orbits. J. Math. Phys. Vol 39, Issue 12, pp. 6372-6383 (1998).
- [LMS1] Lazaroiu CI, McNamee D, Sämann C. Generalized Berezin quantization, Bergman metrics and fuzzy Laplacians. J. High Energy Phys. Issue 9, pp. 1-59 (2008).
- [MaVD] Maes A, Van Daele A. Notes on compact quantum groups. arXiv: math/9803122 (1998).
- [NT1] Neshveyev S, Tuset L. The Martin boundary of a discrete quantum group, J. Reine Angew. Math. Vol 568, pp. 23-70 (2004).
- [Per1] Perelomov AM. Generalized coherent states and applications. Springer (1986).
- [Pims1] Pimsner MV. A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbb{Z} . Fields Inst. Commun. Vol 12, pp. 189-212 (1997).
- [Pop1] Popescu G. Operator theory on noncommutative varieties. Indiana Univ. Math. J. Vol 56, Issue 2, pp. 389-442 (2006).
- [Pop2] Popescu G. Operator theory on noncommutative varieties II. Proc. Amer. Math. Soc. Vol 135, Issue 7, pp. 2151-2164 (2007).
- [Raj1] Rajeev SG. New classical limits of quantum theories. In: Infinite Dimensional Groups and Manifolds. Vol 5, 213 (2004).
- [RCG1] Rawnsley J, Cahen M, Gutt S. Quantization of Kähler manifolds I: geometric interpretation of Berezin's quantization. J. Geom. Phys. Vol 7, Issue 1, pp. 45-62 (1990).
- [RCG2] Rawnsley J, Cahen M, Gutt S. Quantization of Kähler manifolds II. Trans. Amer. Math. Soc. Vol 337, Issue 1 (1993).
- [Rie1] Rieffel M. Deformation quantization for actions of \mathbb{R}^d . Mem. Amer. Math. Soc. Vol 506 (1993).
- [Rie2] Rieffel M. Matrix algebras converge to the sphere for quantum Gromov-Hausdorff distance. Mem. Amer. Math. Soc. Vol 168, Issue 796, pp. 67-91 (2004).
- [RoSt1] Rørdam M, Størmer E (Eds.). Classification of nuclear C^* -algebras. Entropy in operator algebras. Vol 7. Springer (2002).
- [Sain] Sain J. Berezin quantization from ergodic actions of compact quantum groups, and quantum Gromov-Hausdorff distance. PhD thesis, University of California (2009).
- [Schl1] Schlichenmaier M. Berezin-Toeplitz quantization for compact Kähler manifolds. A review of results. Adv. Math. Phys. Article ID 927280, 38 pages (2010).
- [Schl2] Schlichenmaier M. Singular projective varieties and quantization. In: Quantization of Singular Symplectic Quotients. pp. 259-282 (2001).
- [Seg1] Segal G. Lectures on Lie groups. In: Lectures on Lie groups and Lie algebras. Cambridge University Press (1995).
- [Serre1] Serre JP. Géométrie algébrique et géométrie analytique. Ann. Inst. Fourier, Grenoble. Vol 6, pp. 1-42 (1956).
- [ShSo1] Shalit OM, Solel B. Subproduct systems. Doc. Math. Vol 14, pp. 801-868 (2009).
- [Sten1] Stenstrom B. Rings of quotients – An introduction to methods of ring theory. Springer (1975).
- [Timm1] Timmermann T. An invitation to quantum groups and duality: from Hopf algebras to multiplicative unitaries and beyond. Eur. Math. Soc. (2008).

- [Tom1] Tomatsu R. A characterization of right coideals of quotient type and its application to classification of Poisson Boundaries. *Comm. Math. Phys.* Vol 275, Issue 1, pp 271-296 (2007).
- [UnUp1] Unterberger A, Upmeyer H. The Berezin transform and invariant differential operators. *Comm. Math. Phys.* Vol 164, pp. 563-597 (1994).
- [VaVe1] Vaes S, Vennet N. Identification of the Poisson and Martin boundaries of orthogonal discrete quantum groups. *J. Inst. Math. Jussieu.* Issue 7, pp. 391-412 (2008).
- [VVer1] Vaes S, Vergnioux R. The boundary of universal discrete quantum groups, exactness and factoriality. *Duke Math. J.* Vol 140, Issue 1, pp. 35-84 (2006).
- [VaDW] Van Daele A, Wang SZ. Universal quantum groups. *Int. J. Math.* Vol 7, Issue 2, pp. 255-264 (1996).
- [Vas1] Vasselli E. Continuous fields of C^* -algebras arising from extensions of tensor C^* -categories. *J. Funct. Anal.* Vol 199, pp. 122-152 (2003).
- [Vas3] Vasselli E. The C^* -algebra of a vector bundle and fields of Cuntz algebras. *J. Funct. Anal.* 222, Issue 2, pp. 491-502 (2005).
- [Vis2] Viselter A. Cuntz–Pimsner algebras for subproduct systems. *Int. J. Math.* Vol 23, Issue 8, p. 1250081 (2012).
- [Wang1] Wang SZ. Ergodic actions of universal quantum groups on operator algebras. *Comm. Math. Phys.* Vol 203, pp. 481-498 (1999).
- [Wang3] Wang SZ. Structure and isomorphism classification of compact quantum groups $A_u(q)$ and $B_u(q)$. *arXiv:math/9807095v2* (2000).
- [Wa1] Wang X. Balance point and stability of vector bundles over a projective manifold. *Math. Res. Lett.* Vol 9, Issues 2-3, pp. 393-411 (2002).
- [Wor1] Woronowicz SL. Compact matrix pseudogroups. *Comm. Math. Phys.* Vol 111, pp. 613-665 (1987).
- [Wor3] Woronowicz SL. Tannaka–Krein duality for compact matrix pseudogroups. Twisted $SU(n)$ groups, *Inv. Math.* Vol 93, pp. 35-76 (1988).
- [Wor4] Woronowicz SL. Compact quantum groups. *Symétries quantiques, Les Houches 1995*, pp. 845-884 (1998).
- [Yau1] Yau S-T. On the Ricci curvature of a compact Kähler manifold and the complex Monge–Amère equation, I. *Commun. Pure Appl. Math.* Vol 31, pp. 339-411 (1978).
- [Zeld1] Zelditch S. Szegő kernels and a theorem of Tian. *Int. Math. Res. Not.* Issue 6, pp. 317-331 (1998).